

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 October 2003 (30.10.2003)

PCT

(10) International Publication Number  
WO 03/089573 A2

- (51) International Patent Classification<sup>7</sup>: C12N
- (21) International Application Number: PCT/US03/10302
- (22) International Filing Date: 4 April 2003 (04.04.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/370,376 5 April 2002 (05.04.2002) US
- (71) Applicants and  
(72) Inventors: FISHEL, Richard, A. [US/US]; 575 Sprague Road, Penn Valley, PA 19072 (US); YODER, Kristine, E. [US/US]; 575 Sprague Road, Penn Valley, PA 19072 (US).
- (74) Agents: MILLER, Suzanne, E. et al.; Woodcock Washburn LLP, One Liberty Place, 46th Floor, Philadelphia, PA 19103 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**


— of inventorship (Rule 4.17(iv)) for US only

**Published:**

— without international search report and to be republished upon receipt of that report

[Continued on next page]

(54) Title: METHODS OF IDENTIFYING COMPOUNDS THAT MODULATE A DNA REPAIR PATHWAY AND/OR RETROVIRAL INFECTIVITY, THE COMPOUNDS, AND USES THEREOF



(57) Abstract: Retroviruses are RNA viruses that must insert a DNA copy (cDNA) of their genome into the host genome in order to carryout a productive infection. One host cellular pathway that defends against retroviral cDNA integration involves highly conserved proteins of a host DNA repair pathway. These proteins represent novel targets for anti-retroviral drugs. The invention presented herein provides, inter alia, methods of identifying compounds that induce a DNA repair pathway and/or inhibit retroviral cDNA integration into a host genome, compounds thus identified, uses of such compounds, and kits for identifying and testing of the efficacy of compounds in inducing a DNA repair pathway, inhibiting retroviral cDNA integration, and inhibiting retroviral infection.



---

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**METHODS OF IDENTIFYING COMPOUNDS THAT MODULATE  
A DNA REPAIR PATHWAY AND/OR RETROVIRAL INFECTIVITY,  
THE COMPOUNDS, AND USES THEREOF**

**CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims the benefit of U.S. Provisional Application No. 60/370,376, filed April 5, 2002, which is hereby incorporated by reference in its entirety.

**STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH**

[0002] This work was supported in part by a research grant from the National Institutes of Health, grant number GM62556. The United States Government may have certain rights in this invention.

**FIELD OF THE INVENTION**

[0003] The present invention is directed, in part, to methods for inducing a DNA repair pathway, methods for identifying compounds that induce a DNA repair pathway and/or inhibit retroviral infectivity, methods of treating a condition caused by a retroviral infection with compounds that induce a DNA repair pathway and/or inhibit retroviral cDNA integration into the host cell genome, methods for inhibiting a DNA repair pathway and/or increasing retroviral cDNA integration, methods for identifying compounds that inhibit a DNA repair pathway and/or increase retroviral infectivity, and methods of treating a condition by improving gene delivery with compounds that inhibit a DNA repair pathway and/or increase retroviral cDNA integration into the host cell genome.

**BACKGROUND OF THE INVENTION**

[0004] Retroviruses are RNA viruses that must insert a DNA copy (retroviral cDNA) of their genome into the host chromosome in order to carry out a productive infection. Retroviral integration can result in mutagenic inactivation of genes at the sites of cDNA insertion or in aberrant expression of adjacent host genes, both of which can have deleterious consequences for the host organism. Furthermore, retroviruses present considerable risk to human and animal health, as evidenced by the fact that retroviruses cause diseases such as, but not limited to, acquired immune deficiency syndrome (AIDS, caused by human immunodeficiency virus, HIV-1), various animal cancers, feline immunodeficiency virus (FIV), and human adult T-cell leukemia/lymphoma. Retroviruses also have been associated with other common disorders, including, but not limited to, Type I diabetes and multiple sclerosis.

[0005] Recent efforts to combat such retroviral-borne diseases have focused on the identification of inhibitors of retroviral proteins involved in infection. Two mechanisms characterize the mode of infection of retroviruses: reverse transcription and integration (Coffin, J. M., S.H. Hughes, and H.E. Varmus. RETROVIRUSES. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 1997). Both processes are essential for retroviruses to productively infect a cell (Tisdale, M., T. Schulze, B.A. Larder, and K. Moelling. Mutations within the RNase H domain of human immunodeficiency virus type 1 reverse transcriptase abolish virus infectivity. *Journal of General Virology*, 72: 59-66, 1991; LaFemina, R. L., C.L. Schneider, H.L. Robbins, P.O. Callahan, K. LeGrow, E. Roth, W.A. Schleif, and E.A.E. Emini. Requirement of active human immunodeficiency virus type 1 integrase enzyme for productive infection of human T-lymphoid cells. *Journal of Virology*, 66: 7414-7419, 1992; Sakai, H., M. Kawamura, J. Sakuragi, S. Sakuragi, R. Shibata, A. Ishimoto, N. Ono, S. Ueda, and A. Adachi. Integration is essential for efficient gene expression of human immunodeficiency virus type 1. *Journal of Virology*, 67: 1169-1174, 1993; Englund, G., T.S. Theodore, E.O. Freed, A. Engleman, and M.A. Martin. Integration is required for productive infection of monocyte-derived macrophages by human immunodeficiency virus type 1. *Journal of Virology*, 69: 3216-3219, 1995). To date, most drug development programs have focused on inhibition of virally encoded products, including retroviral reverse transcriptases and proteases. However, given the short life cycle of retroviruses and their inherently high rates of genetic change or mutation, such strategies result in the development of drug resistant virus derivatives through alterations of the virally encoded target molecules. Thus, most anti-retroviral drugs that interfere with virally encoded proteins are effective, if at all, for only limited periods of time. Another limitation of



drugs that target retrovirus proteins is that many do not have broad applicability and are highly specific to a particular virus or even a certain strain of a particular virus.

[0006] As an example of the limitations of present retroviral therapies that target retroviral proteins, a current treatment for AIDS, caused by the HIV retrovirus, consists of a cocktail of three or four anti-retroviral drugs termed HAART (highly active anti-retroviral therapy) (Autran, B., G. Carcelain, T.S. Li, C. Blanc, D. Mathez, R. Tubiana, C. Katlama, P. Debre, and J. Leibowitch. Positive effects of combined antiretroviral therapy on CD4+ T cell homeostasis and function in advanced HIV disease. *Science*, 277: 112-116, 1997). The retroviral reverse transcriptase is inhibited by two families of HAART drug components, nucleotide analogs and non-nucleotide inhibitors. The remaining drugs used in HAART are retroviral protease inhibitors, which target another HIV enzyme. However, 78% of new HIV infections are resistant to at least one HAART drug component, and an effective HIV vaccine has not been developed (Richman, D. In: INTERSCIENCE CONFERENCE ON ANTIMICROBIAL AGENTS AND CHEMOTHERAPY, Chicago, IL, 2001; Cohen, J. Debate begins over new vaccine trials. *Science*, 293: 1973, 2001). Furthermore, most of the identified drugs that inhibit the retroviral integrase enzyme of HIV have been unsuccessful in human trials due to lack of specificity or poor bioavailability (Craigie, R. HIV integrase, a brief overview from chemistry to therapeutics. *Journal of Biological Chemistry*, 276: 23213-23216, 2001; Hazuda, D. J., P. Felock, M. Witmer, A. Wolfe, K. Stillmock, J.A. Grobler, A. Espeseth, L. Gabryelski, W. Schleif, C. Blau, and M.D. Miller. Inhibitors of strand transfer that prevent integration and inhibit HIV-1 replication in cells. *Science*, 287: 646-650, 2000). Thus, the development of novel HIV infection and AIDS therapeutics is critical. Also of great importance is the development of an effective HIV vaccine.

[0007] Retroviruses also are used for gene delivery and are likely to play increasingly important roles in gene therapy. Accordingly, methods and compounds that increase retroviral cDNA integration into the host genome, and hence increase gene delivery, are of great importance.

[0008] Thus, an understanding of how retroviruses function and how they can be controlled is of great commercial and medical importance. Such an understanding would allow the development of novel strategies for treating retroviral infection and for improving gene delivery in gene therapy methodologies.

[0009] The present invention elucidates a pathway of DNA repair and its components involved in retrovirus infection and by providing, *inter alia*, methods and assay systems for identifying compounds that inhibit retroviral cDNA integration and/or induce a DNA repair pathway, methods for inducing a DNA repair pathway and/or inhibiting retroviral cDNA integration,

methods of treating a retroviral infection with compounds that induce a DNA repair pathway and/or inhibit retroviral cDNA integration into the host cell genome, methods for inhibiting a DNA repair pathway and/or increasing retroviral cDNA integration, methods for identifying compounds that inhibit a DNA repair pathway and/or increase retroviral infectivity, and methods of treating a condition by improving gene delivery with compounds that inhibit a DNA repair pathway and/or increase retroviral cDNA integration into the host cell genome.

[0010] The stimulation of an intrinsic host defense mechanism as presented herein is a valuable addition to the treatment of HIV, or any other retrovirus, infection. First, it is very difficult or impossible for the retrovirus to mutate in such a way that it evades drug action. Host cell factors are not subject to the highly mutagenic viral replication process, the foundation for development of retroviral drug resistance. Second, since integration is a prerequisite for all retroviruses to be infective, drugs that induce the formation of 1-LTR or 2-LTR circles are effective against a wide spectrum of retrovirus types. Furthermore, little toxicity is associated with this form of treatment since it is an endogenous system (*i.e.*, host cell factors) that is stimulated. The treatment for retroviral infections presented herein is anticipated to be used in combination with other currently available antiviral drugs, for example, as part of HAART.

## SUMMARY OF THE INVENTION

[0011] In one embodiment of the invention, methods for identifying compounds that inhibit retroviral cDNA integration by contacting a cell or cell extract with a non-circularized retroviral cDNA in the presence of a test compound; contacting a cell or cell extract of the same type with a non-circularized retroviral cDNA in the absence of a test compound; and determining whether the amount of retroviral cDNA circularization is increased in the presence of the test compound relative to the level of retroviral cDNA circularization that occurs in the absence of the test compound are provided.

[0012] In another embodiment of the invention, methods for identifying compounds that inhibit retroviral cDNA integration by contacting a cell or cell extract with a non-circularized retroviral cDNA in the presence of a test compound and determining the amount of retroviral cDNA circularization are provided.

[0013] One aspect of the present invention provides methods for identifying compounds that induce a DNA repair pathway in a cell by contacting at least one component of a DNA repair pathway with a non-circularized retroviral cDNA in the presence of a test compound; contacting the component of the DNA repair pathway with a non-circularized retroviral cDNA in the absence of the test compound; and determining whether the amount of retroviral cDNA circularization is increased in the presence of the test compound relative to the amount of

retroviral cDNA circularization that occurs in the absence of the test compound. The methods of the invention may be performed in a cell or in cell extract. Cells that may be employed by the methods of the invention, or from which cell extract may be derived, include, for example, mammalian, including for example human and chicken, yeast, and plant cells. The component of a DNA repair pathway that may be contacted or upregulated, either directly or indirectly, by the test compound includes, but is not limited to, at least one of nucleic acid molecules encoding XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD50, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer; polypeptides encoded thereby; and homologs thereof.

[0014] In some embodiments of the invention, at least one component of a DNA repair pathway exhibits reduced biological activity in the absence of the test compound relative to wild-type biological activity of the component in the absence of the test compound. The component exhibiting reduced biological activity includes, but is not limited to, at least one of nucleic acid molecules encoding XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD50, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer; polypeptides encoded thereby; and homologs thereof.

[0015] In some aspects of the invention, the retroviral cDNA contains at least one marker gene and at least one promoter such that the marker gene is expressed from the promoter upon retroviral cDNA circularization. An increase in retroviral cDNA circularization in the methods of the invention may be detected by an increase in the level of expression of the marker gene or in the level of activity of the polypeptide encoded by the marker gene in the presence of the test compound relative to the level thereof in the absence of the test compound. Examples of marker genes that may be used in the methods of the invention include, but are not limited to, genes encoding green fluorescent protein (GFP), red fluorescent protein (DsRed), alkaline phosphatase (AP),  $\beta$ -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (neor, G418r) dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), lacZ (encoding  $\beta$ -galactosidase), luciferase (luc), or xanthine guanine phosphoribosyltransferase (XGPRT). Examples of promoters that may be used in the methods of the invention include, but are not limited to, promoters derived from adenovirus, SV40, parvoviruses, vaccinia virus, cytomegalovirus, or mammalian genomic DNA, an MSH2 promoter, constitutive promoters including 3-phosphoglycerate kinase and

various other glycolytic enzyme gene promoters, or inducible promoters including the alcohol dehydrogenase-2 promoter or metallothionine promoter.

[0016] Also provided herein are retroviral vectors having a nucleic acid molecule including a promoter and a marker gene that is expressed upon circularization of the nucleic acid molecule. In some embodiments of the invention, the retroviral vector has a nucleic acid sequence of SEQ ID NO:5 or SEQ ID NO:6.

[0017] In some aspects of the invention, compounds that induce a DNA repair pathway and/or inhibit retroviral cDNA integration into the genome of a host cell are provided. In some embodiments of the invention, compounds that prevent retroviral infection of the host cell are provided. In other aspects of the invention, compounds that inhibit a DNA repair pathway and/or increase retroviral cDNA integration are provided.

[0018] Some aspects of the invention are directed to pharmaceutical compositions of the compounds of the invention. Pharmaceutical compositions of the invention, for example for the treatment of a retroviral infection, contain a therapeutically effective amount of at least one compound identified according to the methods of the invention, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

[0019] Additional embodiments of the invention are directed to methods of inducing a DNA repair pathway of a cell by administering at least one compound identified by the methods of the invention to the cell. In some aspects of the invention, the compound inhibits retroviral cDNA integration into the genome of the cell.

[0020] Some embodiments of the invention provide methods of treating a retroviral infection of a patient by administering at least one compound identified by the methods of the invention, or a pharmaceutical composition thereof, to the patient. The patient may be a plant or a mammal, including, but not limited to, avians, felines, canines, bovines, ovines, porcines, equines, rodents, simians, and humans. Examples of retroviral infections that may be treated according to the methods of the invention include, but are not limited to, retroviral infections associated with at least one condition of acquired immune deficiency syndrome (AIDS), human immunodeficiency virus (HIV-1) infection, cancer, human adult T-cell leukemia/lymphoma, FIV, Type I diabetes, and multiple sclerosis.

[0021] One aspect of the present invention provides methods for identifying compounds that inhibit a DNA repair pathway and/or increase retroviral cDNA integration in a cell by contacting at least one component of a DNA repair pathway with a non-circularized retroviral cDNA in the presence of a test compound; contacting the component of the DNA repair pathway with a non-circularized retroviral cDNA in the absence of the test compound; and determining

whether the amount of retroviral cDNA circularization is increased in the presence of the test compound relative to the amount of retroviral cDNA circularization that occurs in the absence of the test compound. The methods of the invention may be performed in a cell or in cell extract. Cells that may be employed by the methods of the invention, or from which cell extract may be derived, include, for example, mammalian, including but not limited to human and chicken, yeast, and plant cells. The component of a DNA repair pathway that may be contacted or upregulated, either directly or indirectly, by the test compound includes, but is not limited to, at least one of nucleic acid molecules encoding XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD50, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer; polypeptides encoded thereby; and homologs thereof.

**[0022]** In some aspects of the invention, the retroviral cDNA contains at least one marker gene and at least one promoter such that the marker gene is expressed from the promoter upon retroviral cDNA circularization. A decrease in retroviral cDNA circularization in the methods of the invention may be detected by a decrease in the level of expression of the marker gene or in the level of activity of the polypeptide encoded by the marker gene in the presence of the test compound relative to the level thereof in the absence of the test compound. Examples of marker genes that may be used in the methods of the invention include, but are not limited to, genes encoding green fluorescent protein (GFP), red fluorescent protein (DsRed), alkaline phosphatase (AP),  $\beta$ -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (neor, G418r) dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), lacZ (encoding  $\beta$ -galactosidase), luciferase (luc), or xanthine guanine phosphoribosyltransferase (XGPRT). Examples of promoters that may be used in the methods of the invention include, but are not limited to, promoters derived from adenovirus, SV40, parvoviruses, vaccinia virus, cytomegalovirus, or mammalian genomic DNA, an MSH2 promoter, constitutive promoters including 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters, or inducible promoters including the alcohol dehydrogenase-2 promoter or metallothionein promoter.

**[0023]** In some aspects of the invention, compounds that inhibit a DNA repair pathway and/or increase retroviral cDNA integration into the genome of a host cell are provided. In some embodiments of the invention, compounds identified according to the methods are provided.

**[0024]** Some aspects of the invention are directed to pharmaceutical compositions of the compounds of the invention. Pharmaceutical compositions of the invention, for example for improving the efficiency of gene delivery in a gene therapy, contain a therapeutically effective

amount of at least one compound identified according to the methods of the invention, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

[0025] Additional embodiments of the invention are directed to methods of inhibiting a DNA repair pathway and/or increasing retroviral cDNA integration of a cell by administering at least one compound identified by the methods of the invention to the cell.

[0026] Additional embodiments of the invention provide methods for increasing the efficiency of gene delivery in a gene therapy by administering a compound of the invention. The patient may be a plant or a mammal, including, but not limited to, avians, felines, canines, bovines, ovines, porcines, equines, rodents, simians, and humans.

[0027] Additional aspects of the invention provide assay systems for identifying compounds that induce a DNA repair pathway. In some aspects of the invention, a cell-free system for identifying a compound that induces a DNA repair pathway containing at least one component of a DNA repair pathway, noncircularized retroviral cDNA having a marker gene that is expressed upon retroviral cDNA circularization, and genomic DNA is provided. Also provided herein are cell-based systems for identifying a compound that induces a DNA repair pathway containing a retrovirus having a marker gene and a cell having at least one component of a DNA repair pathway. In some embodiments of the assay systems, the component of the DNA repair pathway exhibits reduced biological activity relative to wild-type biological activity of the component. In some embodiments of the invention are provided cell-based assay systems for identifying compounds that inhibit retroviral cDNA integration having a cell and a retrovirus containing a circularization marker gene. Also encompassed within the scope of the invention are cell-free assay systems for identifying compounds that inhibit retroviral cDNA integration having host genomic DNA and noncircularized retroviral cDNA having a circularization marker gene.

[0028] Another aspect of the invention is kits containing a retrovirus or retroviral vector of the invention. Such kits may include conventional kit component(s) including but not limited to container(s), label(s), and instructions.

[0029] Other aspects of the invention include methods of screening for a compound which inhibits retroviral infectivity by exposing at least one component of a DNA repair pathway to a test compound; inducing DNA repair; measuring one of an amount of retroviral cDNA circularization wherein the circularization juxtaposes a promoter to a marker gene, and the physical recombination of retroviral cDNA; quantifying expression of the marker gene; inhibiting integration of the retroviral cDNA into a host cell genome; and identifying the compound. Also provided are methods of screening for a compound which inhibits retroviral infectivity by exposing a component of a DNA repair pathway to a test compound; inducing

DNA repair; measuring one of an amount of retroviral cDNA circularization wherein the circularization juxtaposes a promoter to a marker gene, and the physical recombination of retroviral cDNA; measuring an amount of expression of the marker gene which is indicative of an increase in circularization; inhibiting integration of the retroviral cDNA into a host cell genome; and identifying the compound. The component of the DNA repair pathway may be at least one of XPB or XPD but is not limited to the XPB or XPD members of the DNA repair pathway. The component of the DNA repair pathway may be a gene in the DNA repair pathway and the compound which induces DNA repair may upregulate the gene so that DNA repair is induced and retroviral integration is inhibited. The component of the DNA repair pathway also may be a protein in the DNA repair pathway and the compound which induces DNA repair induces an activity or function of the protein so that DNA repair is induced and retroviral integration is inhibited. Additional embodiments of the invention include methods of inhibiting retroviral infectivity in a cell by administering a compound identified to a cell; and inhibiting retrovirus integration into the cell's genome. Also provided are pharmaceutical compositions comprising a compound identified by the screening methods and a pharmaceutically acceptable excipient. A compound that inhibits retroviral integration identified according to the methods herein disclosed. A compound that inhibits retroviral integration identified according to the methods of the invention wherein the compound is a lead compound for further development of a therapeutic agent that causes inhibition of retroviral integration into a host cell's genome.

**[0030]** Additionally provided are methods of inhibiting retroviral infectivity in a subject by administering the test compound identified to a subject and inhibiting retrovirus integration into the genome of the subject. In another embodiment are provided methods of screening for a compound which induces DNA repair in a cell wherein induction of DNA repair inhibits retroviral integration into a host cell's genome by exposing a component of a DNA repair pathway to a test compound; inducing DNA repair; measuring one of an amount of retroviral cDNA circle formation (via homologous recombination or non-homologous end-joining) by quantifying an expression of a marker gene, and the physical recombination of retroviral cDNA; inhibiting integration of the retroviral cDNA into the host cell genome; and identifying the compound. The component of the DNA repair pathway may be at least one of XPB or XPD, but not limited to the XPB or XPD members of the DNA repair pathway. Also encompassed by the invention are methods of inducing DNA repair in a cell wherein induction of DNA repair inhibits retroviral integration into the genome of the cell by administering a test compound identified by a method of the invention to a cell; inducing DNA repair; and inhibiting retrovirus integration into the genome of the cell. Other aspects of the invention include compounds that induce DNA

repair identified according to a method of the invention wherein induction of DNA repair inhibits retroviral integration into a host cell's genome and pharmaceutical compositions of the compound and a pharmaceutically acceptable excipient.

**[0031]** One embodiment of the invention includes methods of inducing DNA repair in a subject by administering a test compound identified to a subject; inducing DNA repair; and inhibiting retrovirus integration into the subject's genome. The compound may induce DNA repair by upregulating a gene in a DNA repair pathway whereby DNA repair is induced and retroviral integration is inhibited or by inducing an activity or function of a protein in a DNA repair pathway whereby DNA repair is induced and retroviral integration is inhibited.

**[0032]** Also provided by the invention are methods of inducing DNA repair in a subject by administering a test compound identified by the methods of the invention to a subject and inducing DNA repair. Compounds that induce DNA repair identified according to methods of the invention may be lead compounds for further development of a therapeutic agent that causes inhibition of retroviral integration into a host cell's genome.

**[0033]** Another aspect of the invention includes methods of screening for a compound which induces DNA repair in a cell wherein induction of DNA repair inhibits retroviral integration into a host cell's genome by exposing a component of a DNA repair pathway to a test compound; inducing DNA repair; measuring one of an amount of retroviral cDNA circle formation (via homologous recombination or non-homologous end-joining) by quantifying an expression of a marker gene, and the physical recombination of retroviral cDNA; and identifying the compound.

**[0034]** The materials, methods, and examples provided herein are illustrative only and are not intended to be limiting. Other features and advantages of the invention will be apparent from the following detailed description and claims.

## **BRIEF DESCRIPTION OF THE DRAWINGS**

**[0035]** **Figures 1A and 1B** illustrate an example of retroviral infection of a host cell. **Figure 1A** shows that HIV infection of a cell begins with the binding of the HIV envelope protein gp120 to the host cell membrane proteins CD4 and either CCR5 or CXCR4. This binding event elicits fusion of the retroviral and cellular membranes, mediated by a second HIV envelope protein gp41. Following membrane fusion, the retroviral capsid core enters the host cell and disassembles in the cytoplasm. HIV reverse transcriptase copies the retroviral genomic RNA into a cDNA molecule. The retroviral cDNA is part of the pre-integration complex (PIC), which includes at least the retroviral proteins integrase, reverse transcriptase, matrix, capsid, and vpr, as well as the host protein HMG I(Y). This complex of protein and nucleic acid enters the host cell nucleus. Retroviral integrase catalyzes the joining of the 3' ends of the retroviral cDNA to the



host genomic DNA. The retroviral cDNA is flanked by five base gaps of host sequence and 5' dinucleotide flaps of HIV sequence. Host DNA repair enzymes finish the integration reaction by repairing the flanking gaps and 5' flaps to generate the provirus. After integration is complete, retroviral and host transcription factors promote the transcription of retroviral mRNAs and genomic RNA. The retroviral mRNAs are translated in the cytoplasm to produce retroviral polyproteins. These polyproteins assemble at the cellular plasma membrane with the retroviral genomic RNA. Immature retroviral particles bud from the cell. After budding, the retroviral enzyme protease cleaves the retroviral polyproteins to yield a mature, infectious virion. **Figure 1B** illustrates that, after the PIC enters the host cell nucleus, integration of the retroviral cDNA will result in a productive infection of the cell. Alternatively, circularization of the retroviral cDNA by one of at least two mechanisms is not productive and will prevent completion of retroviral infection. The host cellular DNA repair mechanism of homologous recombination may generate 1-long terminal repeat (1-LTR) circles. Both ends of the retroviral cDNA have homologous nucleotide sequences, termed long terminal repeats (LTRs). Host DNA repair machinery uses the homologous LTR ends in a recombination reaction to produce 1-LTR circles. A second host cellular DNA repair mechanism, non-homologous end-joining (NHEJ), ligates the ends of the retroviral cDNA to yield 2-long terminal repeat (2-LTR) circles. Neither 1-LTR nor 2-LTR circles can be subsequently converted to retroviral cDNA integration products.

**[0036]** **Figure 2** demonstrates that HIV cDNA integration is controlled by host cell DNA repair. HIV-based vector particles were used to determine relative retroviral cDNA integration efficiency in cell lines varying in DNA repair function. A successful retroviral cDNA integration event is indicated by the expression of green fluorescent protein (GFP) encoded by the HIV vector particles. Cell lines were derived from two patients with mutations of the *XPB* gene (Riou, L., L. Zeng, O. Chevallier-Lagente, A. Stary, O. Nikaido, A. Taieb, G. Weeda, M. Mezzina, and A. Sarasin. The relative expression of mutated *XPB* genes results in xeroderma pigmentosum/Cockayne's syndrome or trichothiodystrophy cellular phenotypes. *Human Molecular Genetics*, 8: 1125-1133, 1999). Three of the cell lines were rescued by addition of an *XPB* transgene. The five cell lines exhibit varying levels of DNA repair requiring *XPB*. The level of *XPB* function is indicated by triangles. The cell lines were transduced with the HIV-based vector particles at relative multiplicities of infection (MOI) of 0, 0.5, and 2, as determined by transduction of 293T human fibroblasts. Following transduction, the cells were fixed and examined by flow cytometry for GFP expression. Cells that did not have vector particles added, 0 MOI, did not express GFP. At both 0.5 MOI and 2 MOI, the percentage of cells expressing GFP (GFP+ cells) was inversely proportional to the level of *XPB* function.

[0037] **Figures 3A and 3B** illustrate one embodiment of a screen for retroviral cDNA circle-formation included within the scope of the invention. **Figure 3A** shows a recombinant retroviral vector constructed to contain a general marker gene (for example, DsRed) driven by a promoter (for example, a cytomegalovirus (CMV) promoter or an MSH2 promoter). Detection of red fluorescence is used as a positive control for retroviral cDNA entry into the host cell nucleus. **Figure 3B** illustrates that the formation of a 1-LTR or 2-LTR circle effectively juxtaposes a second promoter (for example, a CMV promoter or an MSH2 promoter) and a circularization marker gene (for example, GFP) with an intervening LTR (1-LTR or 2-LTR) that is flanked by 5' splice donor and 3' splice acceptor sites. Transcription from this second promoter results in a spliced message that has removed the intervening LTR(s) and will express the circularization marker gene, for example, GFP, and thus be detected, in the case of GFP, as green fluorescence. Because GFP is expressed only upon retroviral cDNA circularization, the level of green fluorescence indicates the efficiency of retroviral cDNA circle-formation versus retroviral cDNA integration into the host cell genome.

[0038] **Figures 4A and 4B** illustrate the nucleotide sequence of the human *XPB* gene (SEQ ID NO:1) and the amino acid sequence of the XPB polypeptide encoded thereby (SEQ ID NO:2), respectively (GenBank Accession No. NM\_000122). **Figures 4C and 4D** provide the nucleotide sequence of the human *XPB* gene (SEQ ID NO:3) and the amino acid sequence of the XPB polypeptide encoded thereby (SEQ ID NO:4), respectively (GenBank Accession No. NM\_000400).

[0039] **Figures 5A-5D** illustrate the nucleotide sequence (SEQ ID NO:5) of one example of the retroviral vector shown in Figure 3, wherein the general marker gene is DsRed, expression of which is controlled by a CMV promoter, and the circularization marker gene is GFP, the expression of which is driven by a CMV promoter upon retroviral cDNA circularization.

[0040] **Figures 6A-6D** illustrate the nucleotide sequence (SEQ ID NO:6) of another example of the retroviral vector shown in Figure 3, wherein the general marker is DsRed, expression of which is controlled by an MSH2 promoter, and the circularization marker gene is GFP, the expression of which is driven by a CMV promoter upon retroviral cDNA circularization.

#### **DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS**

[0041] The reference works, patents, patent applications, and scientific literature that are referred to herein establish the knowledge of those with skill in the art and are hereby incorporated by reference in their entirety to the same extent as if each was specifically and individually indicated to be incorporated by reference. Any conflict between any reference cited herein and the specific teachings of this specification shall be resolved in favor of the latter.

Likewise, any conflict between an art-understood definition of a word or phrase and a definition of the word or phrase as specifically taught in this specification shall be resolved in favor of the latter.

[0042] Standard reference works setting forth the general principles of recombinant DNA technology are known to those of skill in the art (Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, 1998; Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2D ED., Cold Spring Harbor Laboratory Press, Plainview, New York, 1989; Kaufman *et al.*, Eds., HANDBOOK OF MOLECULAR AND CELLULAR METHODS IN BIOLOGY AND MEDICINE, CRC Press, Boca Raton, 1995; McPherson, Ed., DIRECTED MUTAGENESIS: A PRACTICAL APPROACH, IRL Press, Oxford, 1991).

[0043] The present invention relates to the processes whereby retroviruses insert their genetic material into the genome of a eukaryotic host cell in order to carry out a productive infection. More specifically, the present invention relates to highly conserved proteins of the host cell that are required for efficient retroviral cDNA integration. These proteins represent novel targets for anti-retroviral drugs and for drugs for improved gene delivery by retroviruses. Provided herein, *inter alia*, are methods and assay systems that can be used to screen for anti-retroviral compounds and compounds that increase retroviral gene delivery as well as to compare and test similar retroviral assays and drugs *in vivo* and *in vitro*.

[0044] The phrase "DNA repair pathway" as used herein refers to any pathway of a host cell that facilitates repair of the host DNA including but not limited to homologous recombination and non-homologous end-joining. A "component of a DNA repair pathway" refers to any molecule, including but not limited to nucleic acid molecules and polypeptides, that participates in a DNA repair pathway of a host cell. Examples of components of a DNA repair pathway include, but are not limited to, XPA, XPB, XPC, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD50, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and equivalent homologs.

[0045] As used herein, the term "contacting" means bringing together, either directly or indirectly, a compound into physical proximity to a molecule of interest. Contacting may occur, for example, in any number of buffers, salts, solutions, or in a cell or cell extract.

[0046] As used herein, the term "antibody" is meant to refer to complete, intact antibodies, and Fab, Fab', F(ab)2, and other fragments thereof. Complete, intact antibodies include monoclonal antibodies such as murine monoclonal antibodies, chimeric antibodies, and humanized antibodies.

[0047] As used herein, the term "binding" means the physical or chemical interaction between two proteins or compounds or associated proteins or compounds or combinations thereof. Binding includes ionic, non-ionic, Hydrogen bonds, Van der Waals, hydrophobic interactions, *etc.* The physical interaction, the binding, can be either direct or indirect, indirect being through or due to the effects of another protein or compound. Direct binding refers to interactions that do not take place through or due to the effect of another protein or compound but instead are without other substantial chemical intermediates. Binding may be detected in many different manners. As a non-limiting example, the physical binding interaction between two molecules can be detected using a labeled compound. Other methods of detecting binding are well-known to those of skill in the art.

[0048] As used herein, the term "complementary" refers to Watson-Crick basepairing between nucleotide units of a nucleic acid molecule.

[0049] As used herein, the phrase "stringent hybridization conditions" or "stringent conditions" refers to conditions under which an oligonucleotide will specifically hybridize to its target sequence. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the oligonucleotides complementary to the target sequence hybridize to the target sequence at equilibrium. Since the target sequences are generally present in excess, at  $T_m$ , 50% of the hybridizing oligonucleotides are occupied at equilibrium. Typically, stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short oligonucleotides (*e.g.*, 10 to 50 nucleotides) and at least about 60°C for longer oligonucleotides. Stringent conditions may also be achieved with the addition of destabilizing agents, such as formamide.

[0050] The term "marker gene" or "reporter gene" refers to a gene encoding a product that, when expressed, confers a phenotype at the physical, morphologic, or biochemical level on a transformed cell that is easily identifiable, either directly or indirectly, by standard techniques and includes, but is not limited to, green fluorescent protein (GFP), red fluorescent protein (DsRed), alkaline phosphatase (AP),  $\beta$ -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (*neor*, G418r) dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), *lacZ* (encoding  $\beta$ -galactosidase), luciferase (*luc*), and xanthine guanine phosphoribosyltransferase

(XGPRT). As with many of the standard procedures associated with the practice of the invention, skilled artisans will be aware of additional sequences that can serve the function of a marker or reporter. Thus, this list is merely meant to show examples of what can be used and is not meant to limit the invention. The term "general marker" or "general marker gene" as used herein refers to a gene of the retroviral cDNA that is expressed upon integration of the retroviral cDNA into the host genome or upon retroviral cDNA circularization and thus serves as a positive control for retroviral cDNA entry into the host cell nucleus. The term "circularization marker gene" or "circularization marker" refers to a gene of the retroviral cDNA that is expressed only upon circularization of the retroviral cDNA.

[0051] As used herein, the term "promoter" refers to a regulatory element that regulates, controls, or drives expression of a nucleic acid molecule of interest and can be derived from sources such as from adenovirus, SV40, parvoviruses, vaccinia virus, cytomegalovirus, or mammalian genomic DNA. Examples of suitable promoters for mammals include, but are not limited to, CMV and MSH2 promoters. Suitable promoters that can be used in yeast include, but are not limited to, such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine promoter. Again, as with many of the standard procedures associated with the practice of the invention, skilled artisans will be aware of additional promoters that can serve the function of directing the expression of a marker or reporter. Thus, the list is merely meant to show examples of what can be used and is not meant to limit the invention.

[0052] The terms "polypeptide," "peptide," and "protein" are used interchangeably herein.

[0053] As used herein, the term "amino acid" denotes a molecule containing both an amino group and a carboxyl group. In some preferred embodiments, the amino acids are  $\alpha$ -,  $\beta$ -,  $\gamma$ - or  $\delta$ -amino acids, including their stereoisomers and racemates. As used herein the term "L-amino acid" denotes an  $\alpha$ -amino acid having the L configuration around the  $\alpha$ -carbon, that is, a carboxylic acid of general formula  $\text{CH}(\text{COOH})(\text{NH}_2)\text{-(side chain)}$ , having the L-configuration. The term "D-amino acid" similarly denotes a carboxylic acid of general formula  $\text{CH}(\text{COOH})(\text{NH}_2)\text{-(side chain)}$ , having the D-configuration around the  $\alpha$ -carbon. Side chains of L-amino acids include naturally occurring and non-naturally occurring moieties. Non-naturally occurring (*i.e.*, unnatural) amino acid side chains are moieties that are used in place of naturally occurring amino acid side chains in, for example, amino acid analogs. Amino acid substituents may be attached through their carbonyl groups through the oxygen or carbonyl carbon thereof, or through their amino groups, or through functionalities residing on their side chain portions.

[0054] As used herein "polynucleotide" refers to a nucleic acid molecule and includes genomic DNA, cDNA, RNA, mRNA and the like.

[0055] As used herein "antisense oligonucleotide" refers to a nucleic acid molecule that is complementary to at least a portion of a target nucleotide sequence of interest and specifically hybridizes to the target nucleotide sequence under physiological conditions. The term "double stranded RNA" or "dsRNA" as used herein refers to a double stranded RNA molecule capable of RNA interference, including short interfering RNA (siRNA) (see for example, Bass, *Nature*, 411, 428-429 (2001); Elbashir *et al.*, *Nature*, 411, 494-498 (2001)).

[0056] "Synthesized" as used herein refers to polynucleotides produced by purely chemical, as opposed to enzymatic, methods. "Wholly" synthesized DNA sequences are therefore produced entirely by chemical means, and "partially" synthesized DNAs embrace those wherein only portions of the resulting DNA were produced by chemical means.

[0057] "Retroviral cDNA circularization" refers to circle formation, for example 1-LTR or 2-LTR circle formation, of retroviral cDNA.

[0058] "Retroviral cDNA integration" as used herein refers to incorporation of retroviral cDNA into a host cell genomic DNA.

[0059] "Retroviral infection" as used herein refers to the process by which retroviruses propagate within a host cell and includes the steps of reverse transcription of retroviral RNA to retroviral cDNA and integration of retroviral cDNA into the host genome. "Noncircularized retroviral cDNA" or "linear retroviral cDNA" as used herein refers to retroviral cDNA that is not circularized into, for example, a 1-LTR or 2-LTR circle. "Circularized retroviral cDNA" refers to retroviral cDNA that is incapable of integration into a host cell genome and is in the form of a circle, for example, a 1-LTR or 2-LTR circle.

[0060] As used herein, the terms "modulates" or "modifies" means an increase or decrease in the amount, quality, or effect of a particular activity or protein.

[0061] "Inhibitors," "activators," and "modulators" refer to any inhibitory or activating molecules identified using *in vitro* and *in vivo* assays for, *e.g.*, agonists, antagonists, and their homologs, including fragments, variants, and mimetics, as defined herein, that exert substantially the same biological activity as the molecule. "Inhibitors" are compounds that reduce, decrease, block, prevent, delay activation, inactivate, desensitize, or downregulate the biological activity or expression of a molecule or pathway of interest, *e.g.*, antagonists. "Inducers" or "activators" are compounds that increase, induce, stimulate, open, activate, facilitate, enhance activation, sensitize, or upregulate a molecule or pathway of interest, *e.g.*, agonists. In some embodiments of the invention, the level of inhibition or upregulation of the expression or biological activity of

a molecule or pathway of interest refers to a decrease (inhibition or downregulation) or increase (upregulation) of greater than about 50%, 60%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%. The inhibition or upregulation may be direct, *i.e.*, operate on the molecule or pathway of interest itself, or indirect, *i.e.*, operate on a molecule or pathway that affects the molecule or pathway of interest.

[0062] "About" as used herein refers to +/- 10% of the reference value.

[0063] As used herein, "homologous nucleotide sequence" or "homologous amino acid sequence," or variations thereof, refers to sequences characterized by a homology, at the nucleotide level or amino acid level, of at least about 60%, more preferably at least about 70%, more preferably at least about 80%, more preferably at least about 90%, at least about 95%, and most preferably at least about 98% to a reference sequence, or portion or fragment thereof encoding or having a functional domain, for example but not limited to the nucleic acid sequence of SEQ ID NO:1 or SEQ ID NO:3, or a portion of SEQ ID NO:1 or SEQ ID NO:3 which encodes a functional domain of the encoded polypeptide, SEQ ID NO:2 or SEQ ID NO:4, or polypeptides having amino acid sequence SEQ ID NO:2 or SEQ ID NO:4, or fragments thereof having functional domains of the full-length polypeptide. Homologous nucleotide sequences include those sequences coding for homologs, including, for example, isoforms, species variants, allelic variants, and fragments of the protein of interest. Isoforms can be expressed in different tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms can be encoded by different genes. Homologous nucleotide sequences include nucleotide sequences encoding for a species variant of a protein. Homologous nucleotide sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. Homologous amino acid sequences include those amino acid sequences which encode conservative amino acid substitutions in polypeptides having amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4, as well as in polypeptides identified according to the methods of the invention. Percent homology is preferably determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), using the default settings, which uses the algorithm of Smith and Waterman (Smith and Waterman, *Adv. Appl. Math.*, 2: 482-489, 1981). Nucleic acid fragments of the invention have at least about 5, at least about 10, at least about 15, at least about 20, at least about 25, at least about 50, or at least about 100 nucleotides of the reference nucleotide sequence. Preferably the nucleic acid fragments of the invention encode a polypeptide having at least one biological property, or function, that is

substantially similar to a biological property of the polypeptide encoded by the full-length nucleic acid sequence.

[0064] As is well known in the art, because of the degeneracy of the genetic code, there are numerous DNA and RNA molecules that can code for the same polypeptide as that encoded by a nucleotide sequence of interest. The present invention, therefore, contemplates those other DNA and RNA molecules which, on expression, encode a polypeptide encoded by the nucleic acid molecule of interest. DNA and RNA molecules other than those specifically disclosed herein characterized simply by a change in a codon for a particular amino acid, are within the scope of this invention.

[0065] It is to be understood that the present invention includes proteins homologous to, and having at least one biological property, or function, that is substantially similar to a biological property of a reference polypeptide. Preferably, the extent of the biological activity of the biological property is at least 10%, more preferably at least 20%, more preferably at least 30%, more preferably at least 40%, more preferably at least 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 80%, more preferably at least 90%, and most preferably 100% of the activity of the biological property of the reference polypeptide. Such proteins are also called variants. This definition is intended to encompass fragments, isoforms, natural allelic variants, and splice variants. These variant forms may result from, for example, alternative splicing or differential expression in different tissue of the same source organism. The variant forms may be characterized by, for example, amino acid insertion(s), deletion(s), or substitution(s). In this connection, a variant form having an amino acid sequence which has at least about 60%, at least about 70% sequence homology, at least about 80% sequence homology, preferably about 90% sequence homology, more preferably about 95% sequence homology, and most preferably about 98% sequence homology to the reference polypeptide, is included in the present invention. A preferred homologous polypeptide comprises at least one conservative amino acid substitution compared to the reference polypeptide. Amino acid "insertions", "substitutions" or "deletions" are changes to or within an amino acid sequence. The variation allowed in a particular amino acid sequence may be experimentally determined by producing the peptide synthetically or by systematically making insertions, deletions, or substitutions of nucleotides in the nucleic acid sequence using recombinant DNA techniques. Polypeptide fragments of the invention comprise at least about 5, 10, 15, 20, 25, 30, 35, or 40 consecutive amino acids of the reference polypeptide. Preferred polypeptide fragments display antigenic properties unique to, or specific for, the reference polypeptide and its allelic and species



homologs. Fragments of the invention having the desired biological and immunological properties can be prepared by any of the methods well known and routinely practiced in the art.

[0066] Alterations of the naturally occurring amino acid sequence can be accomplished by any of a number of known techniques. For example, mutations can be introduced into the polynucleotide encoding a polypeptide at particular locations by procedures well known to the skilled artisan, such as oligonucleotide-directed mutagenesis, which is described by U.S. Pat. Nos. 4,518,584 and 4,737,462.

[0067] Preferably, a polypeptide homolog of the present invention will exhibit substantially the biological activity of a naturally occurring reference polypeptide. By "exhibit substantially the biological activity of a naturally occurring polypeptide" is meant that variants within the scope of the invention can comprise conservatively substituted sequences, meaning that one or more amino acid residues of a polypeptide are replaced by different residues that do not alter the secondary and/or tertiary structure of the polypeptide. Such substitutions may include the replacement of an amino acid by a residue having similar physicochemical properties, such as substituting one aliphatic residue (Ile, Val, Leu or Ala) for another, or substitution between basic residues Lys and Arg, acidic residues Glu and Asp, amide residues Gln and Asn, hydroxyl residues Ser and Tyr, or aromatic residues Phe and Tyr. Further information regarding making phenotypically silent amino acid exchanges are known in the art (Bowie *et al.*, *Science*, 247: 1306-1310, 1990). Other polypeptide homologs which might retain substantially the biological activities of the reference polypeptide are those where amino acid substitutions have been made in areas outside functional regions of the protein.

[0068] A nucleotide and/or amino acid sequence of a nucleic acid molecule or polypeptide employed in the invention or of a compound identified by the screening method of the invention may be used to search a nucleotide and amino acid sequence databank for regions of similarity using Gapped BLAST (Altschul *et al.*, *Nuc. Acids Res.*, 25: 3389, 1997). Briefly, the BLAST algorithm, which stands for Basic Local Alignment Search Tool is suitable for determining sequence similarity (Altschul *et al.*, *J Mol. Biol.*, 215: 403-410, 1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *J Mol. Biol.*, 215: 403-410, 1990). These initial neighborhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both

directions along each sequence for as far as the cumulative alignment score can be increased. Extension for the word hits in each direction are halted when: 1) the cumulative alignment score falls off by the quantity X from its maximum achieved value; 2) the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or 3) the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (Henikoff *et al.*, *Proc. Natl. Acad. Sci. USA*, 89: 10915-10919, 1992) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands. The BLAST algorithm (Karlin *et al.*, *Proc. Natl. Acad. Sci. USA*, 90: 5873-5787, 1993) and Gapped BLAST perform a statistical analysis of the similarity between two sequences. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a gene or cDNA if the smallest sum probability in comparison of the test nucleic acid to the reference nucleic acid is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

[0069] "Biological activity" as used herein refers to the level of a particular function (for example, enzymatic activity) of a molecule or pathway of interest in a biological system. "Wild-type biological activity" refers to the normal level of function of a molecule or pathway of interest. "Reduced biological activity" refers to a decreased level of function of a molecule or pathway of interest relative to a reference level of biological activity of that molecule or pathway. For example, reduced biological activity may refer to a decreased level of biological activity relative to the wild-type biological activity of a molecule or pathway of interest.

"Increased biological activity" refers to an increased level of function of a molecule or pathway of interest relative to a reference level of biological activity of that molecule or pathway. For example, increased biological activity may refer to an increased level of biological activity relative to the wild-type biological activity of a molecule or pathway of interest.

[0070] As used herein, the term "isolated" nucleic acid molecule refers to a nucleic acid molecule (DNA or RNA) that has been removed from its native environment. Examples of isolated nucleic acid molecules include, but are not limited to, recombinant DNA molecules contained in a vector, recombinant DNA molecules maintained in a heterologous host cell, partially or substantially purified nucleic acid molecules, and synthetic DNA or RNA molecules.

[0071] The term "mimetic" as used herein refers to a compound that is sterically similar to one identified as an inducer of a host DNA repair pathway, provided that the molecule retains

biological activity, *i.e.*, induction of a host DNA repair pathway. Mimetics are structural and functional equivalents to the compounds identified by the present invention that induce a DNA repair pathway.

[0072] The terms "patient" and "subject" are used interchangeably herein and include, but are not limited to, avians, felines, canines, bovines, ovines, porcines, equines, rodents, simians, and humans. "Host cell" includes, for example, a mammalian cell, yeast cell, or plant cell. Mammalian cells of the invention include but are not limited to human and chicken cells (*e.g.*, DT40 cells).

[0073] The term "treatment" as used herein refers to any indicia of success of prevention, treatment, or amelioration of a retroviral infection, or to any indicia of success of improvement of the efficiency of gene delivery in a gene therapy. Treatment of a retroviral infection includes any objective or subjective parameter, such as, but not limited to, abatement, remission, reduction in the number of retroviral particles in a patient, reduction in the number or severity of symptoms or side effects, an increase in the tolerance of the patient to the infection, or slowing of the rate of degeneration or decline of the patient. Treatment of a retroviral infection also includes a prevention of the onset of symptoms in a patient that may be at increased risk of retroviral infection but does not yet experience or exhibit symptoms thereof.

[0074] "Improving efficiency of gene delivery in a gene therapy" refers to any indicia of success of increasing the integration of a gene of a retrovirus or retroviral vector into the host cell genome. "Gene therapy" refers to any treatment method which introduces a gene into a patient for therapeutic effect, for example but not limited to, upregulation or downregulation of an endogenous nucleic acid or polypeptide.

### **Retroviral cDNA integration**

[0075] Some embodiments of the invention disclosed herein inhibit retroviral cDNA integration by stimulating a conserved cellular host defense mechanism, DNA repair. Other embodiments of the invention stimulate retroviral cDNA integration by inhibiting a conserved cellular host defense mechanism. Following reverse transcription, the retrovirus must integrate the cDNA copy of its genome into the host chromosome (Coffin, J. M., S.H. Hughes, and H.E. Varmus. RETROVIRUSES. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 1997; LaFemina, R. L., C.L. Schneider, H.L. Robbins, P.O. Callahan, K. LeGrow, E. Roth, W.A. Schleif, and E.A.E. Emini. Requirement of active human immunodeficiency virus type 1 integrase enzyme for productive infection of human T-lymphoid cells. *Journal of Virology*, 66: 7414-7419, 1992; Sakai, H., M. Kawamura, J. Sakuragi, S. Sakurgai, R. Shibata, A. Ishimoto, N.

Ono, S. Ueda, and A. Adachi. Integration is essential for efficient gene expression of human immunodeficiency virus type 1. *Journal of Virology*, 67: 1169-1174, 1993; Englund, G., T.S. Theodore, E.O. Freed, A. Engleman, and M.A. Martin. Integration is required for productive infection of monocyte-derived macrophages by human immunodeficiency virus type 1. *Journal of Virology*, 69: 3216-3219, 1995). When integrated, the virus is termed a provirus. If a virus is unable to complete the formation of the integrated provirus, it will not be able to continue the infection. The process of retroviral cDNA integration, mediated by the pre-integration complex (PIC), is illustrated in Figure 1A. Host factors that have been shown to influence the integration reaction include, but are not limited to, the high-mobility group protein family (HMGI(Y)), the barrier to autointegration factor (BAF), DNA-dependent protein kinase (DNA-PK), the Ku70/80 heterodimer, XRCC4, and ligase IV (Farnet, C. M., and F.D. Bushman. HIV-1 cDNA integration: requirement of HMG I(Y) protein for function of preintegration complexes *in vitro*. *Cell*, 88: 483-492, 1997; Lee, M. S., and R. Craigie. A previously unidentified host protein protects retroviral DNA from autointegration. *Proceedings of the National Academy of Sciences*, 95: 1528-1533, 1998; Daniel, R., R.A. Katz, A.M. Skalka. A role for DNA-PK in retroviral DNA integration. *Science*, 284, 1999; Li, L., J.M. Olvera, K.E. Yoder, R.S. Mitchell, S.L. Butler, M. Lieber, S.L. Martin, and F.D. Bushman. Role of the non-homologous DNA end-joining pathway in the early steps of retroviral infection. *EMBO Journal*, 20: 3272-3281, 2001). HMGI(Y) and BAF have both been shown to stimulate HIV retroviral cDNA integration *in vitro*. The proteins XRCC4, Ku70/80 heterodimer, and ligase IV catalyze non-homologous end joining (NHEJ) and are able to convert the linear retroviral cDNA to a circular molecule (2-LTR) joined at the long terminal repeat (LTR) sequences (Figure 1B) (Li, L., J.M. Olvera, K.E. Yoder, R.S. Mitchell, S.L. Butler, M. Lieber, S.L. Martin, and F.D. Bushman. Role of the non-homologous DNA end-joining pathway in the early steps of retroviral infection. *EMBO Journal*, 20: 3272-3281, 2001). This 2-LTR circle form of retroviral cDNA is unable to integrate into the host cell genome (Brown, P. O., B. Bowerman, H.E. Varmus, and J.M. Bishop. Retroviral integration: structure of the initial covalent product and its precursor, and a role for the viral IN protein. *Proceedings of the National Academy of Sciences*, 86: 2525-2529, 1989; Engelman, A., G. Englund, J.M. Orenstein, M.A. Martin, and R. Craigie. Multiple effects of mutants in human immunodeficiency virus type 1 integrase on viral replication. *Journal of Virology*, 69: 2729-2736, 1995). An alternative fate for the linear retroviral cDNA is the formation of a 1-LTR circle formed by homologous recombination between the LTRs (Figure 1B).

[0076] The results presented herein demonstrate that stimulation of the formation of 1-LTR and 2-LTR circles of the retroviral cDNA, for example by inducing a DNA repair pathway of a

host cell, inhibits retroviral cDNA integration into the host genome and thus retroviral infectivity. Alternatively, inhibition of 1-LTR and/or 2-LTR circle formation of retroviral cDNA, for example, by inhibiting a DNA repair pathway, increases retroviral cDNA integration into a host cell genome and thus retroviral infectivity.

#### **DNA repair genes control the efficiency of integration**

[0077] During a retroviral infection, nearly all of the linear viral cDNA will either integrate into the host genome or will become 1-LTR or 2-LTR circles (Zennou, V., C. Petit, D. Guetard, U. Nerhbass, L. Montagnier, and P. Charneau HIV-1 genomic nuclear import is mediated by a central DNA flap. *Cell*, 101: 173-185, 2000; Butler, S. L., M.S.T. Hansen, and F.D. Bushman. A quantitative assay for HIV DNA integration *in vivo*. *Nature Medicine*, 7: 631-634, 2001).

Induction of host factors that mediate 1-LTR or 2-LTR circle formation increases the number of 1-LTR or 2-LTR circles, thereby resulting in a decrease in the number of integration events.

Conversely, inhibition or knock-out of host factors that mediate 1-LTR or 2-LTR circle formation decreases retroviral cDNA circularization, thereby resulting in an increase in the number of integration events (Table 1). The invention presented herein describes strategies wherein linear retroviral cDNA molecules that are competent for integration are diverted to the alternative dead-end pathway of 1-LTR or 2-LTR circle formation. The invention also describes strategies for increasing the number of retroviral cDNA integration events by inhibiting 1-LTR or 2-LTR circle formation. Yeast studies suggest that the capacity of this system to control integration is quite large.

[0078] The yeast *Saccharomyces cerevisiae* has been shown to contain a retrovirus-like element family Ty (termed: retrotransposon). The Ty retrotransposon family contains the *gag* and *pol* genes indicative of retroviruses. The *gag* gene encodes all of the structural proteins associated with the virus-like particle. The *pol* gene includes reverse transcriptase, protease and integrase. Polyproteins are translated from the *gag* and *pol* genes and subsequently processed into functional proteins by the protease. Ty lacks an envelope (*env*) gene. Without an *env* gene, Ty particles are unable to bud from the yeast cell and therefore never exist outside the cell. Thus, Ty genomic RNA is transcribed and packaged in the cytoplasm as virus-like particles, that may then be uncoated, reverse transcribed, and integrated into the yeast genome. The lack of an extracellular stage of the life cycle is what defines Ty as a retrotransposon.

[0079] Studies of the Ty retrotransposon in yeast have shown that several yeast cellular DNA repair genes control the efficiency of retroviral cDNA integration. These repair genes include, but are not limited to, *rad25*, *rad3*, *rad50*, *rad51*, *rad52*, *rad54*, and *rad57* (see, for example,

Table 1; Lee, B.-S., C.P. Lichtenstein, B. Faiola, L.A. Rinckel, W. Wysock, M.J. Curcio, and D.J. Garfinkel. Posttranslational inhibition of Ty1 retrotransposition by nucleotide excision repair/transcription factor TFIIH subunits Ssl2p and Rad3p. *Genetics*, 148: 1743-1761, 1998; Rattray, A. J., B.K. Shafer, and D.J. Garfinkel. The *Saccharomyces cerevisiae* DNA recombination and repair functions of the RAD52 epistasis group inhibit Ty1 transposition. *Genetics*, 154: 543-556, 2000). Mutation of these genes leads to great increases in integration efficiency. Conversely, the presence of wild-type DNA repair genes/proteins greatly reduces or prevents the integration reaction.

[0080] Three types of homologous recombination have been identified in eukaryotes that are distinguished by the amount of sequence homology required to induce recombination: microhomology recombination (requiring 1-5 base pairs of homologous sequence between participating parental DNA molecules), short-sequence recombination (requiring 20-300 base pairs of homologous sequence between participating parental DNA molecules), and homologous recombination (requiring >300 base pairs of homologous sequence between participating parental DNA molecules). Microhomology recombination appears to require Rad50p, MRE11p, XRS2(NBS1)p and a DNA ligase (presumed to be XRCC4/Lig4p). Short sequence and homologous recombination appear to require the *rad52*-pathway genes which include, but are not limited to: *rad51*, *rad52*, *rad54*, *rad55*, *rad57*, and *rad59*. In addition, the *rad3* and *rad25* genes also have been found to be part of the short-sequence homologous recombination pathway. All of these recombination pathway genes have human homologs, and all of the pathway types are conserved in human cells.

[0081] Indeed, the same host defense mechanism that inhibits or prevents Ty retrotransposon integration in the yeast *S.cerevisiae* is conserved in mammalian cells, including human. For example, human homologs of the yeast genes *rad25*, *rad3*, *rad1*, *rad2*, *rad14*, *rad50*, *rad51*, *rad52*, *rad54*, *rad57*, *msh2*, and *cdc9* are *XPB*, *XPB*, *XPB*, *XPB*, *XPB*, *hRAD50*, *hRAD51*, *hRAD52*, *hRAD54*, *hRAD57*, *hMSH2*, and *ligase I*, respectively. A number of human genes, including but not limited to *XPB*, *XPB*, *hRAD51*, *hMSH2*, *hRAD51B*, *hRAD51C*, *hRAD51D*, *hXRCC2*, *hXRCC3*, have been identified as components of a human DNA repair pathway involving homologous recombination. The human homologs of Rad25p and Rad3p, *XPB* and *XPB*, respectively, inhibit integration of exogenous DNA (Figure 2). *XPB* and *XPB* have been shown to be helicases that participate in two larger complexes of proteins: the transcription complex TFIIH and the nucleotide excision repair (NER) complex. In humans, mutations in at least one of the seven NER genes (*XPA*, *XPB*, *XPC*, *XPB*, *XPE*, *XPB*, and *XPB*) cause xeroderma pigmentosum (XP), a genetic disease associated with defective NER. NER factors

work together and form multi-protein complexes on damaged DNA (Riou, L., L. Zeng, O. Chevallier-Lagente, A. Sary, O. Nikaido, A. Taieb, G. Weeda, M. Mezzina, and A. Sarasin. The relative expression of mutated XPB genes results in xeroderma pigmentosum/Cockayne's syndrome or trichothiodystrophy cellular phenotypes. *Human Molecular Genetics*, 8: 1125-1133, 1999).

**[0082]** The present invention shows that the DNA helicases XPB and XPD participate in the transformation of the linear retroviral cDNA to circularized retroviral cDNA, for example 1-LTR circles. The formation of 1-LTR circles is controlled by homologous recombination between the direct repeat LTRs of the retroviral cDNA. The level of retroviral cDNA integration inhibition is inversely proportional to the level of XPB repair activity *in vivo* (Figure 2).

**[0083]** A second host cellular DNA repair mechanism, non-homologous end-joining (NHEJ), ligates the ends of the retroviral cDNA to yield 2-long terminal repeat (2-LTR) circles. The proteins DNA-PK, Ku70/80 heterodimer, XRCC4, ligase IV, hMRE11, hRAD50, and XRS2 (NBS1) participate in NHEJ. Members of the NHEJ pathway, including Ku70/80 heterodimer, ligase IV, and XRCC4, have been shown to convert the linear retroviral cDNA to a circular molecule (2-LTR) joined at the long terminal repeat (LTR) sequences (Figure 1B).

#### **DNA repair pathway and anti-retroviral action**

**[0084]** Inhibition of at least one component of a DNA repair pathway increases retroviral cDNA integration. Stimulation of at least one component of a DNA repair pathway decreases retroviral cDNA integration.

**[0085]** In some aspects of the present invention, genes and/or proteins within a DNA repair pathway are induced, that is, DNA repair is stimulated in order to inhibit retroviral cDNA integration. In some embodiments of the present invention the expression of a gene in a DNA repair pathway is upregulated, thereby increasing the production of at least one component of a DNA repair pathway. In some embodiments of the present invention, the biological activity or function of a protein involved in DNA repair is induced by a compound that interacts directly or indirectly with at least one component of a DNA repair pathway.

**[0086]** In some aspects of the present invention, genes and/or proteins within a DNA repair pathway are inhibited in order to increase retroviral cDNA integration. In some embodiments of the present invention the expression of a gene in a DNA repair pathway is downregulated, thereby decreasing the production of at least one component of a DNA repair pathway. In some embodiments of the present invention, the activity or function of a protein involved in DNA

repair is decreased by a compound that interacts directly or indirectly with at least one protein of a DNA repair pathway.

### Screening for compounds

[0087] The present invention provides methods for identifying compounds that modulate retroviral cDNA integration into a host genome. In some aspects of the invention, components of a DNA repair pathway have uses in the screening methods to detect molecules that specifically induce or inhibit components of a DNA repair pathway or bind the components of a DNA repair pathway to enhance or reduce their activity. In one embodiment, such assays are performed to screen molecules for utility as anti-retroviral drugs or lead compounds for drug development.

[0088] Methods of screening for compounds that modulate retroviral cDNA integration into the host genome include contacting a cell or cell extract with a non-circularized retroviral cDNA in the presence of a test compound and measuring the retroviral cDNA circularization that occurs. The amount of retroviral cDNA circularization that occurs in the presence of the test compound(s) may be compared with the retroviral cDNA circularization that occurs in comparable reaction medium that is not treated with the test compound(s). Compounds that increase retroviral cDNA integration cause a decrease of retroviral cDNA circularization as compared to the control in the absence of the test compound(s). Compounds that decrease retroviral cDNA integration cause an increase of retroviral cDNA circularization as compared to the control in the absence of the test compound(s).

[0089] Methods of screening for compounds that induce DNA repair include the steps of contacting one or more test compounds with one or more components of a DNA repair pathway of an organism of interest (which organism can be one of many different species, including, but not limited to, avians, felines, canines, bovines, ovines, porcines, equines, rodents, simians, and humans) in a suitable reaction medium and testing for compound/component interaction, *e.g.* by assessing the activity of the DNA repair pathway, or component thereof, and comparing that activity with the activity in comparable reaction medium that is not treated with the test compound(s). A difference in the activity between the treated and untreated samples is indicative of a modulating effect of the relevant test compound(s). Prior to being screened for the ability actually to affect or modulate DNA repair, test compounds may be screened for their ability to physically interact with a component of a DNA repair pathway. This may, for example, be used as a coarse screen prior to testing a compound for actual ability to modulate biological activity.



[0090] The components of a DNA repair pathway employed in the screening assay may be provided in a cell to be exposed to the test compound. Alternatively the assay may be performed on an *in vitro* DNA repair system that measures the accuracy and efficiency of joining together DNA strand breaks that have been created by treating intact DNA with restriction endonucleases, chemicals, radiation, or a recombinant retrovirus.

[0091] The activation of a DNA repair pathway leads to the protection of host DNA from degradation and thus protection from retroviral cDNA integration. Activation of a DNA repair pathway may be caused by DNA double-strand breaks (DSBs), single strand gaps in the DNA double helix, or by other disruptions to the DNA double-helix. These structures exist at the ends of retroviral cDNA and occur as intermediates in the retroviral cDNA integration process. Assays for DNA repair, retrovirus or retroviral cDNA, intermediates in retroviral cDNA integration, or synthetic preparations of DNA that mimic any of these may be provided.

[0092] Methods of the invention identify compounds that modulate DNA repair and/or retroviral cDNA integration by their ability to modulate retroviral cDNA circle (1-LTR or 2-LTR) formation. Induction of DNA repair or inhibition of retroviral cDNA integration by the test compound is verified by an increase in retroviral cDNA circle-formation. Inhibition of DNA repair or stimulation of retroviral cDNA integration by the test compound is verified by a decrease in retroviral cDNA circle-formation. Retroviral cDNA circle-formation is scored using standard genetic, biochemical, cellular, or histological techniques. For example, but not meant to limit the invention, a retroviral vector is designed such that the short-sequence homologous recombination that leads to the formation of the 1-LTR circles or non-homologous end-joining that leads to the formation of 2-LTR circles results in the juxtaposition of a promoter and a circularization marker gene, such as, but not limited to, green fluorescent protein (GFP) (Figure 3). Proximity of the promoter to the marker gene results in expression of the marker gene, such as GFP, thereby allowing for the direct measurement of the expressed marker gene by cellular or biochemical techniques. The present invention also contemplates assaying for the ability of a test compound to affect the biological activity of a component of a DNA repair pathway. Thus, for example, compounds may be screened for their ability to affect DNA-PK phosphorylation, *etc.*

[0093] Screening of organic or peptide libraries with expressed recombinant protein components of a DNA repair pathway is useful for identification of therapeutic molecules that modulate the activity of a DNA repair pathway. In one embodiment screening is carried out to select for compounds that stimulate DNA repair as determined by the induction of 1-LTR or 2-

LTR formation. In another embodiment, screening is performed to select for compounds that inhibit DNA repair as determined by the inhibition of 1-LTR or 2-LTR formation.

[0094] Diversity libraries, such as random or combinatorial peptide or non-peptide libraries are also screened for molecules that specifically stimulate or inhibit DNA repair. Many libraries are known in the art that can be used, such as, but not limited to, chemically synthesized libraries, recombinant (*e.g.*, phage display libraries), and *in vitro* translation-based libraries. By way of examples of non-peptide libraries, a benzodiazepine library can be used. Peptide libraries can also be used. Another example of a library that can be used is one in which the amide functionalities in peptides have been permethylated to generate a chemically transformed combinatorial library. These methods are well known to those of skill in the art and can be found in standard molecular technique references.

[0095] Screening the libraries can be accomplished by any of a variety of commonly known methods.

#### **The test system**

[0096] Host cells for the methods of the invention are preferably eukaryotic cells. Given the ease of manipulation of yeast, an assay according to the present invention may involve applying test compounds to a yeast system. Mammalian cells, including but not limited to human cells and chicken cells (*e.g.*, DT40 cells), and plant cells also may be used in the methods of the invention.

[0097] For therapeutic purposes, a DNA repair pathway, or one or more components (or subunits) thereof, may be employed in the assay. The DNA repair pathway, or components thereof, may be, for example but not limited to, avian, feline, bovine, ovine, porcine, equine, rodent, simian, or human. In view of the high conservation between DNA repair components in different eukaryotes, similar results will be obtained using the compounds in mammalian, *e.g.* human, systems. In other words, a compound identified as being able to induce DNA repair in yeast will be able to induce DNA repair in other eukaryotes. A further approach is to employ standard recombinant technology techniques to generate yeast cells that express one or more components or subunits of a DNA repair pathway of another eukaryote, *e.g.* human. A plant DNA repair pathway, or one or more components thereof or cells comprising the components, may also be used in an assay according to the present invention to test for a compound(s) useful in modulating retrotransposon or retroelement activity in plants.

[0098] Alternatively, the system for screening for compounds in the methods of the invention may be cell-free, *e.g.*, in a cell extract.

**Compounds identified by the screening methods**

[0099] A compound that tests positive in an assay according to the present invention, *i.e.*, is found to inhibit retroviral cDNA integration and/or stimulate DNA repair or, alternatively, is found to inhibit DNA repair and/or increase retroviral cDNA integration, may be peptide or non-peptide in nature. As used herein, the term “compound” means any identifiable chemical or molecule, including, but not limited to, small molecule, peptide, protein, sugar, nucleotide, or nucleic acid, and such compound can be natural or synthetic. Such compounds may include, for example, antibodies, antisense oligonucleotides, and small molecules. A “compound” identified by a screening method of the invention includes the compound so identified, in addition to homologs and mimetics thereof having the same functional effect on DNA repair and/or retroviral cDNA integration.

**Antisense and siRNA**

[0100] Compounds that inhibit DNA repair identified according to the methods of the invention include antisense oligonucleotides and small interfering RNA (siRNA) molecules to a component of a DNA repair pathway.

[0101] Antisense oligonucleotides are administered to cells or cell extract to disrupt at least one component of a DNA repair pathway. The antisense oligonucleotides hybridize to polynucleotides encoding a component of a DNA repair pathway. Both full-length and polynucleotide fragments are suitable for use as antisense oligonucleotides. “Antisense oligonucleotide fragments” of the invention include, but are not limited to oligonucleotides that specifically hybridize to DNA or RNA encoding a component of a DNA repair pathway (as determined by a sequence comparison of oligonucleotides encoding a component of a DNA repair pathway to oligonucleotides encoding other known polypeptides). Examples of antisense oligonucleotides of the invention include but are not limited to antisense oligonucleotides that hybridize to SEQ ID NO:1 or SEQ ID NO:3. Identification of sequences that are substantially unique to DNA repair component-encoding oligonucleotides can be ascertained by analysis of any publicly available sequence database and/or with any commercially available sequence comparison programs. Antisense molecules may be generated by any means including, but not limited to chemical synthesis, expression in an *in vitro* transcription reaction, expression in a transformed cell comprising a vector that may be transcribed to produce antisense molecules, restriction digestion and isolation, the polymerase chain reaction, and the like.

[0102] Those of skill in the art recognize that the antisense oligonucleotides that inhibit the expression and/or biological activity of a component of a DNA repair pathway may be predicted

using any genes encoding a component of a DNA repair pathway. Specifically, antisense nucleic acid molecules comprise a sequence complementary to at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 100, 250 or 500 nucleotides or an entire DNA repair gene sequence. Preferably, the antisense oligonucleotides comprise a sequence complementary to about 15 consecutive nucleotides of the coding strand of the DNA repair component-encoding sequence.

[0103] In one embodiment, an antisense nucleic acid molecule is antisense to a “coding region” of the coding strand of a nucleotide sequence encoding a DNA repair pathway component protein. The coding strand may also include regulatory regions of the DNA repair pathway component sequence. The term “coding region” refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues. In another embodiment, the antisense nucleic acid molecule is antisense to a “noncoding region” of the coding strand of a nucleotide sequence encoding a DNA repair protein. The term “noncoding region” refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions (UTR)).

[0104] Antisense oligonucleotides may be directed to regulatory regions of a nucleotide sequence encoding a DNA repair protein, or mRNA corresponding thereto, including, but not limited to, the initiation codon, TATA box, enhancer sequences, and the like. Given the coding strand sequences provided herein, antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of a DNA repair component mRNA, but also may be an oligonucleotide that is antisense to only a portion of the coding or noncoding region of the mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of an mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length.

[0105] Another means to inhibit the activity of a DNA repair pathway component according to the invention is via RNA interference (RNAi) (*see e.g.*, Elbashir *et al.*, *Nature*, 411:494-498 (2001); Elbashir *et al.*, *Genes Development*, 15:188-200 (2001)). RNAi is the process of sequence-specific, post-transcriptional gene silencing, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene (*e.g.*, is homologous in sequence to the sequence of a DNA repair pathway component, for example but not limited to the sequence as set forth in SEQ ID NO:1 or SEQ ID NO:3). siRNA-mediated silencing is thought to occur post-transcriptionally and/or transcriptionally. For example, siRNA duplexes may mediate post-

transcriptional gene silencing by reconstitution of siRNA-protein complexes (siRNPs), which guide mRNA recognition and targeted cleavage.

[0106] Accordingly, another form of a DNA repair pathway inhibitory compound of the invention is a short interfering RNA (siRNA) directed against a DNA repair pathway component-encoding sequence. Exemplary siRNAs are siRNA duplexes (for example, 10-25, preferably 20, 21, 22, 23, 24, or 25 residues in length) having a sequence homologous or identical to a fragment of the XPB sequence set forth as SEQ ID NO:1 or the XPD sequence of SEQ ID NO:3, and having a symmetric 2-nucleotide 3'-overhang. The 2-nucleotide 3' overhang is preferably composed of (2'-deoxy) thymidine because it reduces costs of RNA synthesis and may enhance nuclease resistance of siRNAs in the cell culture medium and within transfected cells. Substitution of uridine by thymidine in the 3' overhang is also well tolerated in mammalian cells, and the sequence of the overhang appears not to contribute to target recognition.

### Antibodies

[0107] Also comprehended by the present invention are antibodies (*e.g.*, monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies, bifunctional/bispecific antibodies, humanized antibodies, human antibodies, and complementary determining region (CDR) grafted antibodies, including compounds which include CDR sequences which specifically recognize a polypeptide of the invention) specific for components of a DNA repair pathway or fragments thereof. Preferred antibodies of the invention are human antibodies that are produced and identified according to methods described in WO93/11236, published June 20, 1993. Antibody fragments, including Fab, Fab', F(ab')<sub>2</sub>, and Fv, are also provided by the invention. The term "specific for," when used to describe antibodies of the invention, indicates that the variable regions of the antibodies of the invention recognize and bind a component of a DNA repair pathway exclusively (*i.e.*, are able to distinguish the component from other known molecules by virtue of measurable differences in binding affinity, despite the possible existence of localized sequence identity, homology, or similarity). It will be understood that specific antibodies may also interact with other proteins (for example, *S. aureus* protein A or other antibodies in ELISA techniques) through interactions with sequences outside the variable region of the antibodies, and, in particular, in the constant region of the molecule. Screening assays to determine binding specificity of an antibody of the invention are well known and routinely practiced in the art. For a comprehensive discussion of such assays, see Harlow et al. (Eds.), ANTIBODIES A LABORATORY MANUAL; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988), Chapter 6. Antibodies that recognize and bind fragments of a component of a DNA

repair pathway of the invention are also contemplated, provided that the antibodies are specific for the component of the DNA repair pathway. Antibodies of the invention can be produced using any method well known and routinely practiced in the art.

[0108] The invention provides an antibody that is specific for a component of a DNA repair pathway or an epitope thereof. Examples of antibodies of the invention include but are not limited to antibodies to the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO:4, or epitopes thereof. Antibody specificity is described in greater detail below. Cross-reactive antibodies are not antibodies that are "specific" for a component of a DNA repair pathway. The determination of whether an antibody is specific or is cross-reactive with another molecule is made using any of several assays, such as Western blotting assays, that are well known in the art.

[0109] In one preferred variation, the invention provides monoclonal antibodies. Hybridomas that produce such antibodies also are intended as aspects of the invention. In yet another variation, the invention provides a humanized antibody. Humanized antibodies are useful for *in vivo* therapeutic indications.

[0110] In another variation, the invention provides a cell-free composition comprising polyclonal antibodies, wherein at least one of the antibodies is an antibody of the invention specific for a component of a DNA repair pathway. Antisera isolated from an animal is an exemplary composition, as is a composition comprising an antibody fraction of an antisera that has been resuspended in water or in another diluent, excipient, or carrier.

[0111] In still another related embodiment, the invention provides an anti-idiotypic antibody specific for an antibody that is specific for a component of a DNA repair pathway.

[0112] It is well known that antibodies contain relatively small antigen binding domains that can be isolated chemically or by recombinant techniques. Such domains are useful DNA repair pathway component-binding molecules themselves, and also may be reintroduced into human antibodies, or fused to toxins or other polypeptides. Thus, in still another embodiment, the invention provides a polypeptide comprising a fragment of a DNA repair pathway component-specific antibody, wherein the fragment and the polypeptide bind to the component of a DNA repair pathway. By way of non-limiting example, the invention provides polypeptides that are single-chain antibodies and CDR (complementary determining region)-grafted antibodies.

[0113] Non-human antibodies may be humanized by any of the methods known in the art. In one method, the non-human CDRs are inserted into a human antibody or consensus antibody framework sequence. Further changes can then be introduced into the antibody framework to modulate affinity or immunogenicity.

[0114] Antibodies of the invention are useful for, *e.g.*, therapeutic purposes (by modulating activity of a component of a DNA repair pathway).

### Mimetics

[0115] Mimetics or mimics of compounds identified herein (sterically similar compounds formulated to mimic the key portions of the structure) may be designed for pharmaceutical use. Mimetics may be used in the same manner as the compounds identified by the present invention that stimulate DNA repair and hence are also functional equivalents. The generation of a structural-functional equivalent may be achieved by the techniques of modeling and chemical design known to those of skill in the art. It will be understood that all such sterically similar constructs fall within the scope of the present invention.

[0116] The designing of mimetics to a known pharmaceutically active compound is a known approach to the development of pharmaceuticals based on a "lead" compound. This is desirable where the active compound is difficult or expensive to synthesize, or where it is unsuitable for a particular method of administration, *e.g.* peptides are unsuitable active agents for oral compositions as they tend to be quickly degraded by proteases in the alimentary canal. There are several steps commonly taken in the design of a mimetic from a compound that induces DNA repair. First, the particular parts of the compound that are critical and/or important in determining its DNA repair-inducing properties are determined. In the case of a polypeptide, this can be done by systematically varying the amino acid residues in the peptide, *e.g.* by substituting each residue in turn. Alanine scans of peptides are commonly used to refine such peptide motifs.

[0117] Once the active region of the compound has been identified, its structure is modeled according to its physical properties, *e.g.* stereochemistry, bonding, size and/or charge, using data from a range of sources, such as, but not limited to, spectroscopic techniques, X-ray diffraction data, and NMR. Computational analysis, similarity mapping (which models the charge and/or volume of the active region, rather than the bonding between atoms), and other techniques known to those of skill in the art can be used in this modeling process.

[0118] In a variant of this approach, the three-dimensional structure of the compound that induces DNA repair and the active region of the target component of a DNA repair pathway are modeled. This can be especially useful where either or both of these compounds change conformation on binding.

[0119] A template molecule is then selected onto which chemical groups that mimic the compound that induces DNA repair can be grafted. The template molecule and the chemical

groups grafted onto it can conveniently be selected so that the mimetic is easy to synthesize, is pharmacologically acceptable, and does not degrade *in vivo*, while retaining the biological activity of the lead compound. Alternatively, where the mimetic is peptide-based, further stability can be achieved by cyclizing the peptide, thereby increasing its rigidity. The mimetic or mimetics found by this approach can then be screened by the methods of the present invention to see whether they have the ability to induce DNA repair. Further optimization or modification can then be carried out to arrive at one or more final mimetics for *in vivo* or clinical testing.

### **Pharmaceutical compositions**

[0120] Following identification of a compound that induces DNA repair and/inhibits retroviral cDNA integration or, alternatively, inhibits DNA repair and/or stimulates retroviral cDNA integration, the compound may be manufactured and/or used in preparation of a pharmaceutical composition. These are administered to patients, including, but are not limited to, avians, felines, canines, bovines, ovines, porcines, equines, rodents, simians, and humans.

[0121] Thus, the present invention extends, in various aspects, not only to compounds identified in accordance with the methods disclosed herein but also pharmaceutical compositions, drugs, or other compositions comprising such a compound; methods comprising administration of such a composition to a patient, *e.g.* for treatment (which includes prophylactic treatment) of a retroviral disorder or for improving the efficiency of gene delivery in a gene therapy; uses of such a compound in the manufacture of a composition for administration to a patient; and methods of making a composition comprising admixing such a compound with a pharmaceutically acceptable excipient, vehicle or carrier, and optionally other ingredients.

[0122] The pharmaceutical compositions of the invention comprise a therapeutically effective amount of a compound identified according to the methods disclosed herein, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient.

[0123] The compounds of the invention can be formulated as neutral or salt forms.

Pharmaceutically acceptable salts include those formed with free amino groups such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, *etc.*, and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, *etc.*

[0124] Pharmaceutically acceptable carriers include but are not limited to saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The carrier and composition can be sterile. The formulation should suit the mode of administration.



[0125] The composition, if desired, can also contain minor amounts of wetting or emulsifying agents or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, *etc.*

[0126] In one embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for oral (*e.g.*, tablets, granules, syrups) or non-oral (*e.g.*, ointments, injections) administration to the subject. Various delivery systems are known and can be used to administer a compound that induces DNA repair and/or inhibits retroviral cDNA integration, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, expression by recombinant cells, receptor-mediated endocytosis, construction of a therapeutic nucleic acid as part of a retroviral or other vector, *etc.* Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, topical, and oral routes.

[0127] The compounds of the invention may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, *etc.*), and may be administered together with other biologically active agents, for example in HAART therapy. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir.

[0128] In a specific embodiment, it may be desirable to administer the pharmaceutical compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery; topical application, *e.g.*, in conjunction with a wound dressing after surgery; by injection; by means of a catheter; by means of a suppository; or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers.

[0129] The composition can be administered in unit dosage form and may be prepared by any of the methods well known in the pharmaceutical art, for example, as described in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Publishing Co., Easton, PA). The amount of the compound of the invention that induces DNA repair and/or inhibits retroviral cDNA integration or, alternatively, that inhibits DNA repair and/or increase retroviral cDNA integration that is

effective in the treatment of a particular disorder or condition will depend on factors including but not limited to the chemical characteristics of the compounds employed, the route of administration, the age, body weight, and symptoms of a patient, the nature of the disorder or condition, and can be determined by standard clinical techniques. Typically therapy is initiated at low levels of the compound and is increased until the desired therapeutic effect is achieved. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges. Suitable dosage ranges for intravenous administration are generally about 20-500 micrograms of active compound per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.01 pg/kg body weight to 1 mg/kg body weight. Suppositories generally contain active ingredient in the range of 0.5% to 10% by weight; oral formulations preferably contain 10% to 95% active ingredient. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[0130] Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry-lyophilized powder or water-free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline.

[0131] Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

### **Treatment Methods**

[0132] The invention provides methods of treatment of retroviral infections by administration to a subject or patient of an effective amount of a compound that induces DNA repair and/or inhibits retroviral cDNA integration into the host genome. In some aspects of the invention, the compounds or pharmaceutical compositions of the invention are administered to a patient having an increased risk of or having a retroviral infection. The patient may be, for example, avian, feline, canine, bovine, ovine, porcine, equine, rodent, simian, or human. The retroviral infection may be associated with at least one of acquired immune deficiency syndrome (AIDS), human immunodeficiency virus (HIV) infection, cancer, human adult T-cell leukemia, lymphoma, FIV, Type I diabetes, and multiple sclerosis.

[0133] The invention also provides methods of treatment, for example, by improving gene delivery, by administering to a patient or subject an effective amount of a compound that increases retroviral cDNA integration and/or inhibits DNA repair. The patient may be, for example, avian, feline, canine, bovine, ovine, porcine, equine, rodent, simian, or human.

#### Kits of Retroviruses Having a Circularization Marker Gene

[0134] A kit of the invention comprises a carrier means being compartmentalized to receive in close confinement one or more container means such as vials, tubes, and the like, each of the container means comprising an element to be used in the methods of the invention. For example, one of the container means may comprise the retrovirus or retroviral vector of the invention having a circularization marker gene. The kit may also have one or more conventional kit components, including, but not limited to, instructions, test tubes, Eppendorf<sup>TM</sup> tubes, labels, reagents helpful for quantification of marker gene expression, *etc.*

**TABLE 1. DNA Repair Pathway Component Knockouts Increase Integration**

| <i>Saccharomyces cerevisiae</i><br>gene | Fold-increase<br>Ty<br>Transposition <sup>1</sup> | Human<br>gene | Fold-increase<br>HIV<br>Integration <sup>2</sup> |
|---|---|---------------|--|
| <i>rad25</i>                            | 2-1125 <sup>3</sup>                               | <i>XPB</i>    | 2.5  |
| <i>rad3</i>                             | 17-41 <sup>3</sup>                                | <i>XPB</i>    | 2  |
| <i>rad1</i>                             | 1   |               |  |
| <i>rad2</i>                             | 1   |               |  |
|   |   | <i>XPA</i>    | 1  |
| <i>rad50</i>                            | 12  |               |  |
| <i>rad51</i>                            | 11  |               |  |
| <i>rad52</i>                            | 24  |               |  |
| <i>rad54</i>                            | 5   |               |  |
| <i>rad57</i>                            | 21  |               |  |
| <i>msh2</i>                             | 1   | <i>hMSH2</i>  | 1  |
| <i>cdc9</i>                             | 38  |               |  |

<sup>1</sup>Continuous transposition/integration throughout the growth and analysis of cells.

<sup>2</sup>Single-round infection-generated increase in integration.

<sup>3</sup>Significant Stain Background effects

**Additional references**

1. Temin, H. M., E. Keshet, and S.K. Weller. Correlation of transient accumulation of linear unintegrated viral DNA and transient cell killing by avian leukosis and reticuloendotheliosis viruses. *Cold Spring Harbor Symposium on Quantitative Biology*, 44: 773-778, 1980.
2. Pang, S., Y. Koyanagi, S. Miles, C. Wiley, H.V. Vintners, and I.S.Y. Chen. High levels of unintegrated HIV-1 DNA in brain tissue of AIDS dementia patients. *Science*, 343: 85-89, 1990.
3. Bergeron, L., and J. Sodroski. Dissociation of unintegrated viral DNA accumulation from single-cell lysis induced by human immunodeficiency virus type 1. *Journal of Virology*, 66: 5777-5787, 1992.
4. Berg, D. E., and M.M. Howe. *Mobile DNA*. Washington, DC: American Society for Microbiology, 1989.
5. Curcio, M. J., A.M. Hedge, J.D. Boeke, and D.J. Garfinkel. Ty RNA levels determine the spectrum of retrotransposition events that activate gene expression in *Saccharomyces cerevisiae*. *Molecular General Genetics*, 220: 213-221, 1990.
6. Curcio, M. J., and D.J. Garfinkel. Single-step selection for Ty1 element retrotransposition. *Proceedings of the National Academy of Sciences*, 88: 936-940, 1991.
7. Slupianek, A., C. Schmutte, G. Tomblin, M. Nieborowska-Skorska, G. Hoser, M.O. Nowicki, A.J. Pierce, R. Fishel, and T. Skorski. BCR/ABL regulates mammalian RecA homologs, resulting in drug resistance. *Molecular Cell*, 8: 795-806, 2001.

**What is claimed is:**

1. A method of screening for a compound which induces a DNA repair pathway of a cell, comprising:
  - a) contacting at least one component of a DNA repair pathway with a non-circularized retroviral cDNA in the presence of a test compound;
  - b) contacting said at least one component of a DNA repair pathway with a non-circularized retroviral cDNA in the absence of said test compound; and
  - c) determining whether the amount of retroviral cDNA circularization is increased in the presence of said test compound relative to the amount of retroviral cDNA circularization in the absence of said test compound.
2. The method according to claim 1, wherein said component contacted with the test compound is a nucleic acid molecule encoding a polypeptide selected from the group consisting of XPA, XPB, XPC, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.
3. The method according to claim 2, wherein said nucleic acid molecule encodes XPB or XPD.
4. The method according to claim 3, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
5. The method according to claim 3, wherein said nucleic acid molecule encoding XPB has a nucleotide sequence of SEQ ID NO:1 and wherein said nucleic acid molecule encoding XPD has a nucleotide sequence of SEQ ID NO:3.
6. The method according to claim 1, wherein said component contacted with the test compound is a polypeptide selected from the group consisting of XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.

7. The method according to claim 6, wherein said polypeptide is XPB or XPD.
8. The method according to claim 7, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
9. The method according to claim 1, wherein at least one component of said DNA repair pathway in the absence of said test compound exhibits reduced biological activity relative to wild-type biological activity of said component.
10. The method according to claim 9, wherein said component exhibiting reduced biological activity is a nucleic acid molecule encoding a polypeptide selected from the group consisting of XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.
11. The method according to claim 10, wherein said nucleic acid molecule encodes XPB or XPD.
12. The method according to claim 11, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
13. The method according to claim 11, wherein said nucleic acid molecule encoding XPB has a nucleotide sequence of SEQ ID NO:1 and wherein said nucleic acid molecule encoding XPD has a nucleotide sequence of SEQ ID NO:3.
14. The method according to claim 9, wherein said component exhibiting reduced biological activity is a polypeptide selected from the group consisting of XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.
15. The method according to claim 14, wherein said polypeptide is XPB or XPD.

16. The method according to claim 15, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
17. The method according to claim 1, wherein said test compound directly or indirectly upregulates the expression of at least one component of a DNA repair pathway.
18. The method according to claim 17, wherein said upregulated component of a DNA repair pathway is a nucleic acid molecule encoding a polypeptide selected from the group consisting of XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.
19. The method according to claim 18, wherein said nucleic acid molecule encodes XPB or XPD.
20. The method according to claim 19, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
21. The method according to claim 19, wherein said nucleic acid molecule encoding XPB has a nucleotide sequence of SEQ ID NO:1 and wherein said nucleic acid molecule encoding XPD has a nucleotide sequence of SEQ ID NO:3.
22. The method according to claim 1, wherein said test compound directly or indirectly upregulates the biological activity of at least one component of a DNA repair pathway.
23. The method according to claim 22, wherein said upregulated component of a DNA repair pathway is a polypeptide selected from the group consisting of XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.
24. The method according to claim 22, wherein said polypeptide is XPB or XPD.

25. The method according to claim 24, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
26. The method according to claim 1, wherein said retroviral cDNA comprises at least one marker gene and at least one promoter and wherein said marker gene is expressed from said promoter upon retroviral cDNA circularization.
27. The method according to claim 26, wherein said increase in retroviral cDNA circularization is detected by an increase in the level of expression of said marker gene in the presence of said test compound relative to the level of expression of said marker gene in the absence of said test compound.
28. The method according to claim 26, wherein said increase in retroviral cDNA circularization is detected by an increase in the level of activity of the polypeptide expressed from said marker gene in the presence of said test compound relative to the level of activity of the polypeptide expressed from said marker gene in the absence of said test compound.
29. The method according to claim 27, wherein said marker gene encodes green fluorescent protein (GFP), red fluorescent protein (DsRed), alkaline phosphatase (AP),  $\beta$ -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (neor, G418r) dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), lacZ (encoding  $\beta$ -galactosidase), luciferase (luc), or xanthine guanine phosphoribosyltransferase (XGPRT).
30. The method according to claim 26, wherein said promoter is an adenovirus promoter, an SV40 promoter, a parvovirus promoter, a vaccinia virus promoter, a cytomegalovirus promoter, an MSH2 promoter, or a mammalian genomic DNA promoter.
31. The method according to claim 26, wherein said promoter is a 3-phosphoglycerate kinase gene promoter, an alcohol dehydrogenase-2 promoter, or a metallothionine promoter.
32. The method according to claim 1, wherein steps (a) and (b) occur in a cell or in cell extract.



33. The method according to claim 32, wherein said cell is a mammalian or yeast cell.
34. The method according to claim 1, wherein said compound inhibits retroviral cDNA integration into the genome of a cell.
35. The method of claim 34, wherein said compound prevents retroviral infection.
36. A compound that induces a DNA repair pathway of a cell identified according to the method of claim 1.
37. A pharmaceutical composition for the treatment of a retroviral infection comprising a therapeutically effective amount of at least one compound identified according to the method of claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.
38. A method of inducing a DNA repair pathway of a cell comprising administering at least one compound identified according to the method of claim 1 to said cell.
39. The method according to claim 38, wherein said compound inhibits retroviral cDNA integration into the genome of said cell.
40. A method of treating a retroviral infection of a patient comprising administering at least one compound identified according to the method of claim 1 to said patient.
41. The method according to claim 40, wherein said patient is a mammal.
42. The method according to claim 41, wherein said mammal is avian, feline, canine, bovine, ovine, porcine, equine, rodent, simian, or human.
43. The method according to claim 42, wherein said mammal is a human.
44. The method according to claim 40, wherein said retroviral infection is associated with at least one condition selected from the group consisting of acquired immune deficiency syndrome (AIDS), human immunodeficiency virus (HIV) infection, cancer, human adult T-cell leukemia, lymphoma, feline immunodeficiency virus (FIV), Type I diabetes, and multiple sclerosis.

45. The method according to claim 45, wherein said retroviral infection is HIV infection or AIDS.
46. A kit for identifying a compound that induces a DNA repair pathway comprising a retrovirus or retroviral vector having a marker gene that is expressed upon retroviral cDNA circularization.
47. The kit according to claim 46, further comprising at least one conventional kit component.
48. Use of a compound identified according to the method of claim 1 in the manufacture of a pharmaceutical composition for the treatment of a retroviral infection.
49. A method of identifying a compound that inhibits retroviral cDNA integration into a host genome comprising:
- a) contacting a first cell or cell extract with a non-circularized retroviral cDNA in the presence of a test compound;
  - b) contacting a second cell or cell extract with a non-circularized retroviral cDNA in the absence of said test compound, wherein said first and said second cell or cell extract are of the same cell type; and
  - c) determining whether the amount of retroviral cDNA circularization is increased in the presence of said test compound relative to the amount of retroviral cDNA circularization in the absence of said test compound.
50. The method according to claim 49, wherein said retroviral cDNA comprises at least one marker gene and at least one promoter and wherein said marker gene is expressed from said promoter upon retroviral cDNA circularization.
51. The method according to claim 50, wherein said increase in retroviral cDNA circularization is detected by an increase in the level of expression of said marker gene in the presence of said test compound relative to the level of expression of said marker gene in the absence of said test compound.

52. The method according to claim 50, wherein said increase in retroviral cDNA circularization is detected by an increase in the level of activity of the polypeptide expressed from said marker gene in the presence of said test compound relative to the level of activity of the polypeptide expressed from said marker gene in the absence of said test compound.
53. The method according to claim 50, wherein said marker gene encodes green fluorescent protein (GFP), red fluorescent protein (DsRed), alkaline phosphatase (AP),  $\beta$ -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (neor, G418r) dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), lacZ (encoding  $\beta$ -galactosidase), luciferase (luc), or xanthine guanine phosphoribosyltransferase (XGPRT).
54. The method according to claim 50, wherein said promoter is an adenovirus promoter, an SV40 promoter, a parvovirus promoter, a vaccinia virus promoter, a cytomegalovirus promoter, an MSH2 promoter, or a mammalian genomic DNA promoter.
55. The method according to claim 50, wherein said promoter is a 3-phosphoglycerate kinase gene promoter, an alcohol dehydrogenase-2 promoter, or a metallothionine promoter.
56. The method according to claim 49, wherein said cell type is mammalian or yeast.
57. A compound that inhibits retroviral cDNA integration into a host cell genome identified according to the method of claim 49.
58. A pharmaceutical composition for the treatment of a retroviral infection comprising a therapeutically effective amount of at least one compound identified according to the method of claim 49, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.
59. A method of inhibiting retroviral cDNA integration into a host cell genome by administering a compound identified according to the method of claim 49 to said cell.
60. A method of treating a retroviral infection of a patient comprising administering at least one compound identified according to the method of claim 49 to said patient.

61. The method according to claim 60, wherein said patient is a mammal.
62. The method according to claim 61, wherein said mammal is avian, feline, canine, bovine, ovine, porcine, equine, rodent, simian, or human.
63. The method according to claim 62, wherein said mammal is a human.
64. The method according to claim 60, wherein said retroviral infection is associated with at least one condition selected from the group consisting of acquired immune deficiency syndrome (AIDS), human immunodeficiency virus (HIV) infection, cancer, human adult T-cell leukemia, lymphoma, feline immunodeficiency virus (FIV), Type I diabetes, and multiple sclerosis.
65. The method according to claim 64, wherein said retroviral infection is HIV infection or AIDS.
66. Use of a compound identified according to the method of claim 49 in the manufacture of a pharmaceutical composition for the treatment of a retroviral infection.
67. A kit for identifying a compound that inhibits retroviral cDNA integration into a host genome comprising a retrovirus or retroviral vector having a marker gene that is expressed upon retroviral cDNA circularization.
68. The kit according to claim 67, further comprising at least one conventional kit component.
69. A retroviral vector comprising a nucleic acid molecule having promoter and a marker gene that is expressed upon circularization of said nucleic acid molecule.
70. The retroviral vector of claim 69, wherein said marker gene encodes green fluorescent protein (GFP), red fluorescent protein (DsRed), alkaline phosphatase (AP),  $\beta$ -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (neor, G418r) dihydrofolate reductase (DHFR), hygromycin-B-

phosphotransferase (HPH), thymidine kinase (TK), lacZ (encoding  $\beta$ -galactosidase), luciferase (luc), or xanthine guanine phosphoribosyltransferase (XGPRT).

71. The retroviral vector of claim 69, wherein said promoter is an adenovirus promoter, an SV40 promoter, a parvovirus promoter, a vaccinia virus promoter, a cytomegalovirus promoter, an MSH2 promoter, or a mammalian genomic DNA promoter.
72. The retroviral vector of claim 69, wherein said promoter is a 3-phosphoglycerate kinase gene promoter, an alcohol dehydrogenase-2 promoter, or a metallothionine promoter.
73. The retroviral vector of claim 69 comprising the nucleotide sequence of SEQ ID NO:5 or SEQ ID NO:6.
74. A method of screening for a compound which induces a DNA repair pathway of a cell, comprising:
- a) contacting at least one component of a DNA repair pathway with a non-circularized retroviral cDNA in the presence of a test compound; and
  - b) determining the amount of retroviral cDNA circularization.
75. A method of identifying a compound that inhibits retroviral cDNA integration into a host genome comprising:
- a) contacting a cell or cell extract with a non-circularized retroviral cDNA in the presence of a test compound; and
  - b) determining the amount of retroviral cDNA circularization.
76. A method of screening for a compound which inhibits a DNA repair pathway of a cell, comprising:
- a) contacting at least one component of a DNA repair pathway with a non-circularized retroviral cDNA in the presence of a test compound;
  - b) contacting said at least one component of a DNA repair pathway with a non-circularized retroviral cDNA in the absence of said test compound; and
  - c) determining whether the amount of retroviral cDNA circularization is decreased in the presence of said test compound relative to the amount of retroviral cDNA circularization in the absence of said test compound.

77. The method according to claim 76, wherein said component contacted with the test compound is a nucleic acid molecule encoding a polypeptide selected from the group consisting of XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.
78. The method according to claim 77, wherein said nucleic acid molecule encodes XPB or XPD.
79. The method according to claim 78, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
80. The method according to claim 78, wherein said nucleic acid molecule encoding XPB has a nucleotide sequence of SEQ ID NO:1 and wherein said nucleic acid molecule encoding XPD has a nucleotide sequence of SEQ ID NO:3.
81. The method according to claim 76, wherein said component contacted with the test compound is a polypeptide selected from the group consisting of XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.
82. The method according to claim 81, wherein said polypeptide is XPB or XPD.
83. The method according to claim 82, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
84. The method according to claim 76, wherein said test compound directly or indirectly downregulates the expression of at least one component of a DNA repair pathway.
85. The method according to claim 84, wherein said downregulated component of a DNA repair pathway is a nucleic acid molecule encoding a polypeptide selected from the group consisting of XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3,

XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.

86. The method according to claim 85, wherein said nucleic acid molecule encodes XPB or XPD.
87. The method according to claim 86, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
88. The method according to claim 86, wherein said nucleic acid molecule encoding XPB has a nucleotide sequence of SEQ ID NO:1 and wherein said nucleic acid molecule encoding XPD has a nucleotide sequence of SEQ ID NO:3.
89. The method according to claim 76, wherein said test compound directly or indirectly downregulates the biological activity of at least one component of a DNA repair pathway.
90. The method according to claim 89, wherein said downregulated component of a DNA repair pathway is a polypeptide selected from the group consisting of XPA, XPB, XPC, XPD, XPE, XPF, XPG, RAD50, RAD52, RAD54, RAD57, RAD59, MSH2, CDC9, hRAD51, hRAD51B, hRAD51C, hRAD51D, hXRCC2, hXRCC3, XRCC4, ligase IV, hMRE11, XRS2 (NBS1), DNA-PK, and Ku70/80 heterodimer, and homologs thereof.
91. The method according to claim 89, wherein said polypeptide is XPB or XPD.
92. The method according to claim 91, wherein said XPB has an amino acid sequence of SEQ ID NO:2 and wherein said XPD has an amino acid sequence of SEQ ID NO:4.
93. The method according to claim 76, wherein said retroviral cDNA comprises at least one marker gene and at least one promoter and wherein said marker gene is expressed from said promoter upon retroviral cDNA circularization.
94. The method according to claim 93, wherein said decrease in retroviral cDNA circularization is detected by a decrease in the level of expression of said marker gene in the

presence of said test compound relative to the level of expression of said marker gene in the absence of said test compound.

95. The method according to claim 93, wherein said decrease in retroviral cDNA circularization is detected by a decrease in the level of activity of the polypeptide expressed from said marker gene in the presence of said test compound relative to the level of activity of the polypeptide expressed from said marker gene in the absence of said test compound.
96. The method according to claim 93, wherein said marker gene encodes green fluorescent protein (GFP), red fluorescent protein (DsRed), alkaline phosphatase (AP),  $\beta$ -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (neor, G418r) dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), lacZ (encoding  $\beta$ -galactosidase), luciferase (luc), or xanthine guanine phosphoribosyltransferase (XGPRT).
97. The method according to claim 93, wherein said promoter is an adenovirus promoter, an SV40 promoter, a parvovirus promoter, a vaccinia virus promoter, a cytomegalovirus promoter, an MSH2 promoter, or a mammalian genomic DNA promoter.
98. The method according to claim 93, wherein said promoter is a 3-phosphoglycerate kinase gene promoter, an alcohol dehydrogenase-2 promoter, or a metallothionine promoter.
99. The method according to claim 76, wherein steps (a) and (b) occur in a cell or in cell extract.
100. The method according to claim 99, wherein said cell is a mammalian or yeast cell.
101. The method according to claim 76, wherein said compound increases retroviral cDNA integration into the genome of a cell.
102. A compound that inhibits a DNA repair pathway of a cell identified according to the method of claim 76.



103. A pharmaceutical composition for increasing efficiency of gene delivery in a gene therapy comprising a therapeutically effective amount of at least one compound identified according to the method of claim 76, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.
104. A method of inhibiting a DNA repair pathway of a cell comprising administering at least one compound identified according to the method of claim 76 to said cell.
105. The method according to claim 104, wherein said compound increases retroviral cDNA integration into the genome of said cell.
106. A method of improving efficiency of gene delivery in a gene therapy of a patient comprising administering at least one compound identified according to the method of claim 76 to said patient.
107. The method according to claim 106, wherein said patient is a mammal.
108. The method according to claim 107, wherein said mammal is avian, feline, canine, bovine, ovine, porcine, equine, rodent, simian, or human.
109. The method according to claim 108, wherein said mammal is a human.
110. A kit for identifying a compound that inhibits a DNA repair pathway comprising a retrovirus or retroviral vector having a marker gene that is expressed upon retroviral cDNA circularization.
111. The kit according to claim 110, further comprising at least one conventional kit component.
112. Use of a compound identified according to the method of claim 76 in the manufacture of a pharmaceutical composition for increasing the efficiency of gene delivery in a gene therapy.
113. A method of identifying a compound that increases retroviral cDNA integration into a host genome comprising:

- a) contacting a first cell or cell extract with a non-circularized retroviral cDNA in the presence of a test compound;
- b) contacting a second cell or cell extract with a non-circularized retroviral cDNA in the absence of said test compound, wherein said first and said second cell or cell extract are of the same cell type; and
- c) determining whether the amount of retroviral cDNA circularization is decreased in the presence of said test compound relative to the amount of retroviral cDNA circularization in the absence of said test compound.

114. The method according to claim 113, wherein said retroviral cDNA comprises at least one marker gene and at least one promoter and wherein said marker gene is expressed from said promoter upon retroviral cDNA circularization.

115. The method according to claim 113, wherein said decrease in retroviral cDNA circularization is detected by a decrease in the level of expression of said marker gene in the presence of said test compound relative to the level of expression of said marker gene in the absence of said test compound.

116. The method according to claim 114, wherein said decrease in retroviral cDNA circularization is detected by a decrease in the level of activity of the polypeptide expressed from said marker gene in the presence of said test compound relative to the level of activity of the polypeptide expressed from said marker gene in the absence of said test compound.

117. The method according to claim 114, wherein said marker gene encodes green fluorescent protein (GFP), red fluorescent protein (DsRed), alkaline phosphatase (AP),  $\beta$ -lactamase, chloramphenicol acetyltransferase (CAT), adenosine deaminase (ADA), aminoglycoside phosphotransferase (neor, G418r) dihydrofolate reductase (DHFR), hygromycin-B-phosphotransferase (HPH), thymidine kinase (TK), lacZ (encoding  $\beta$ -galactosidase), luciferase (luc), or xanthine guanine phosphoribosyltransferase (XGPRT).

118. The method according to claim 114, wherein said promoter is an adenovirus promoter, an SV40 promoter, a parvovirus promoter, a vaccinia virus promoter, a cytomegalovirus promoter, an MSH2 promoter, or a mammalian genomic DNA promoter.

119. The method according to claim 114, wherein said promoter is a 3-phosphoglycerate kinase gene promoter, an alcohol dehydrogenase-2 promoter, or a metallothionine promoter.
120. The method according to claim 113, wherein said cell type is mammalian or yeast.
121. A compound that increases retroviral cDNA integration into a host cell genome identified according to the method of claim 113.
122. A pharmaceutical composition for the increasing the efficiency of gene delivery in a gene therapy comprising a therapeutically effective amount of at least one compound identified according to the method of claim 113, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.
123. A method of increasing retroviral cDNA integration into a host cell genome by administering a compound identified according to the method of claim 113 to said cell.
124. A method of improving the efficiency of gene delivery of a gene therapy of a patient comprising administering at least one compound identified according to the method of claim 113 to said patient.
125. The method according to claim 124, wherein said patient is a mammal.
126. The method according to claim 125, wherein said mammal is avian, feline, bovine, ovine, porcine, equine, rodent, simian, or human.
127. The method according to claim 126, wherein said mammal is a human.
128. Use of a compound identified according to the method of claim 113 in the manufacture of a pharmaceutical composition for improving the efficiency of gene delivery in a gene therapy.
129. A kit for identifying a compound that increases retroviral cDNA integration into a host genome comprising a retrovirus or retroviral vector having a marker gene that is expressed upon retroviral cDNA circularization.

130. The kit according to claim 129, further comprising at least one conventional kit component.

131. A method of screening for a compound which inhibits a DNA repair pathway of a cell, comprising:

- a) contacting at least one component of a DNA repair pathway with a non-circularized retroviral cDNA in the presence of a test compound; and
- b) determining the amount of retroviral cDNA circularization.

132. A method of identifying a compound that increases retroviral cDNA integration into a host genome comprising:

- a) contacting a cell or cell extract with a non-circularized retroviral cDNA in the presence of a test compound; and
- b) determining the amount of retroviral cDNA circularization.





**Figure 1A****Figure 1B**


Figure 2  
HIV Integration Is Controlled By DNA Repair



**Figure 3A**  
***Retroviral Provirus***



**Figure 3B**  
***Retroviral 1-LTR Circle***



***Retroviral 2-LTR Circle***




Figure 4A

1 GGGAGCTTCCGGATTGAGCCGGAAGTCCCCCAGAGCGGATGCCGCGGCGGGCCTGTGGG  
61 AGCGGGGTCATCTTCTCTCTGCTGCTGTAGCTGCCATGGGCAAAAGAGACCGAGCGGACC  
121 GCGACAAGAAGAAATCCAGGAAGCGGCACTATGAGGATGAAGAGGATGATGAAGAGGACG  
181 CCCCAGGGAACGACCCTCAGGAAGCGGTTCCCTCGGCGGCGGGGAAGCAGGTGGATGAGT  
241 CAGGCACCAAAGTGGATGAATATGGAGCCAAGGACTACAGGCTGCAAATGCCGCTGAAGG  
301 ACGACCACACCTCCAGGCCCTCTGGGTGGCTCCCGATGGCCATATCTTCTTGAAGCCT  
361 TCTCTCCAGTTTACAAATATGCCCAAGACTTCTTGGTGGCTATTGCAGAGCCAGTGTGCC  
421 GACCAACCCATGTGCATGAGTACAACTAACTGCCTACTCCTTGTATGCAGCTGTCAGCG  
481 TTGGGCTGCAAACCAGTGACATCACCGAGTACCTCAGGAAGCTCAGCAAGACTGGAGTCC  
541 CTGATGGAATTATGCAGTTTATTAAGTTGTGTACTGTCAGCTATGGAAAAGTCAAGCTGG  
601 TCTTGAAGCACAAACAGATACTTCGTTGAAAGTTGCCACCCTGATGTAATCCAGCATCTTC  
661 TCCAGGACCCCGTGATCCGAGAATGCCGCTTAAGAACTCTGAAGGGGAGGCCACTGAGC  
721 TCATCACAGAGACTTTCACAAGCAAATCTGCCATTTCTAAGACTGCTGAAAGCAGTGGTG  
781 GGCCCTCCACTTCCCGAGTGACAGATCCACAGGGTAAATCTGACATCCCCATGGACCTGT  
841 TTGACTTCTATGAGCAAATGGACAAGGATGAAGAAGAAGAAGAAGAGACACAGACAGTGT  
901 CTTTTGAAGTCAAGCAGGAAATGATTGAGGAACTCCAGAAACGTTGCATCCACCTGGAGT  
961 ACCCTCTGTTGGCAGAATATGACTTCCGGAATGATTCTGTCAACCCTGATATCAACATTG  
1021 ACCTAAAGCCCACAGCTGTCCCTCAGACCCTATCAGGAGAAGAGCTTGCGAAAGATGTTTG  
1081 GAAACGGGCGTGACGTTCCGGGGGTCATTGTTCTTCCCTGCGGTGCTGGAAAGTCCCTGG  
1141 TTGGTGTGACTGCTGCATGCACTGTCAGAAAACGCTGTCTGGTGCTGGGCAACTCAGCTG  
1201 TTTCTGTGGAGCAGTGGAAAGCCCAGTTCAGATGTGGTCCACCATTGACGACAGCCAGA  
1261 TCTGCCGGTTCACCTCCGATGCCAAGGACAAGCCCATCGGCTGCTCCGTTGCCATTAGCA  
1321 CCTACTCCATGCTGGGCCACACCACCAAAAGGTCTGGGAGGCCGAGCGAGTCATGGAGT  
1381 GGCTCAAGACCCAGGAGTGGGGCCTCATGATCCTGGATGAAGTGCACACCATAACCAGCCA  
1441 AGATGTTCCGAAGGGTGCTCACCATCGTGCAGGCCCACTGTAAGCTGGGTTTGAAGTGC  
1501 CCCTCGTCCGCGAAGATGACAAAATTGTGGATTTAAATTTTCTGATTGGGCCTAAGCTCT  
1561 ACGAAGCCAACCTGGATGGAGCTGCAGAATAATGGCTACATCGCCAAAGTCCAGTGTGCTG  
1621 AGGTCTGGTGCCCTATGTCTCCTGAATTTTACCGGGAATATGTGGCAATCAAAACCAAGA  
1681 AACGAATCTTGCTGTACACCATGAACCCCAACAAATTTAGAGCTTGCCAGTTTCTGATCA  
1741 AGTTTCATGAAAGGAGGAATGACAAGATTATTGTCTTTGCTGACAATGTGTTTGCCCTAA  
1801 AGGAATATGCCATTCGACTGAACAAACCCTATATCTACGGACCTACGTCTCAGGGGGAAA  
1861 GGATGCAAATTTCTCAGAATTTCAAGCACAAACCCCAAAATTAACACCATCTTCATATCCA  
1921 AGGTAGGTGACACTTCGTTTGATCTGCCGGAAGCAAATGTCCTCATTGAGATCTCATCCC  
1981 ATGGTGGCTCCAGGCGTCAGGAAGCCCAAGGCTAGGGCGGGTGCTTCGAGCTAAAAAAG  
2041 GGATGGTTGCAGAAGAGTACAATGCCTTTTTTCTACTCACTGGTATCCCAGGACACACAGG  
2101 AAATGGCTTACTCAACCAAGCGGCAGAGATTCTTGGTAGATCAAGGTTATAGCTTCAAGG  
2161 TGATCACGAAACTCGCTGGCATGGAGGAGGAAGACTTGGCGTTTTTCGACAAAAGAAGAGC  
2221 AACAGCAGCTCTTACAGAAAGTCCTGGCAGCCACTGACCTGGATGCCGAGGAGGAGGTGG  
2281 TGGCTGGGGAATTTGGCTCCAGATCCAGCCAGGCATCTCGGCGCTTTGGCACCATGAGTT  
2341 CTATGTCTGGGGCCGACGACACTGTGTACATGGAGTACCACTCATCGCGGAGCAAGGCGC  
2401 CCAGCAAACATGTACACCCGCTCTTCAAGCGCTTTAGGAAATGATGCTTAGGCAGGGTAC  
2461 TTCGTTCAAGACCGGCGCTTGGCACCCCTTGTGGAAAGGGATTTTCAGCATAACATTTTC  
2521 CTTCCACCTCTTTGACCTTCCCTCCAGCGTTGGCCAAATTTGTGCTGAGGAAGATGCATCA  
2581 AGGGCTTGGCTGTGCCTTCATAGGTCATCTAGGGTTTTATAAAGGAGGAGGAGACAATAT  
2641 TTTTTCAAACTTTTTGGGGAGTGGGGTCATTTCTGTATATAAAAAATGTTAATATTTAAG  
2701 GTGTATTTATGTTACCGTTCTGAATAAACAGAATGGACCATTGAACCAGTA



**Figure 4B**

MGKRDRADRDKKKSRKRHYEDEEDDEEDAPGNDPQEAVPSAAGKQVDESGTKVDEYGAKDYRLQ  
MPLKDDHTSRPLWVAPDGHIFLEAFSPVYKYAQDFLVAIAEPVCRPTHVHEYKLTAYSLYAASV  
VGLQTS DITEYLRKLSKTGVDPDGIMQFIKLCTVSYGKVKLVLKHNRYFVESCHPDV IQHLLQDP  
VIRECRLRNSEGEATELITETFTSKSAISKTAESSGGPSTSRVTDPQGKSDIPMDLFD FYEQMD  
KDEEEEEETQTVSFEVKQEMIEELQKRCIHLEYPLLAEYDFRND SVNPDINIDLKPTAVLRPYQ  
EKSLRKMFGNGRARS GVIVLPCGAGKSLVGVTA ACTVRKRCLVLGNSAVSVEQWKAQFKMWSTI  
DDSQICRFTSDAKDKPIGCSVAISTYSMLGHTTKRSWEAERVMEWLKTQEWGLMILDEVHTIPA  
KMFRRVLTIVQAHCKLGLTATLVREDDKIVDLN FLIGPKLYEANWMELQNNGYIAKVQCAEVWC  
PMSPEFYREYVAIKTKKRILLYTMNPNKFRACQFLIKFHERRNDKIIVFADNVFALKEYAIRLN  
KPYIYGPTSQGERMQILQNFKHNP KINTIFISKVGDTSFDLPEANVLIQISSHGGSSRRQEAQRL  
GRVLRAKKGMVAEEYN AFFYSLVSQDTQEMAYSTKRQRFLVDQGY SFKVITKLAGMEEEDLAFS  
TKEEQQLLQKVLAATDLDAEEEVVAGEFGSRSSQASRRFGTMSSMSGADDTVYMEYHSSRSKA  
PSKHVHPLFKRFRK

Figure 4C

1 ATGAAGCTCAACGTGGACGGGCTCCTGGTCTACTTCCCGTACGACTACATCTACCCCGAG  
61 CAGTTCTCCTACATGCGGGAGCTCAAACGCACGCTGGACGCCAAGGGTCATGGAGTCCTG  
121 GAGATGCCCTCAGGCACCGGGAAGACAGTATCCCTGTTGGCCCTGATCATGGCATAACAG  
181 AGAGCATATCCGCTGGAGGTGACCAAACCTCATCTACTGCTCAAGAACTGTGCCAGAGATT  
241 GAGAAGGTGATTGAAGAGCTTCGAAAGTTGCTCAACTTCTATGAGAAGCAGGAGGGCGAG  
301 AAGCTGCCGTTTCTGGGACTGGCTCTGAGCTCCCGCAAAAACCTTGTGTATTACCCCTGAG  
361 GTGACACCCCTGCGCTTTGGGAAGGACGTGATGGGAAATGCCACAGCCTCACAGCCTCC  
421 TATGTGCGGGCGCAGTACCAGCATGACACCAGCCTGCCCCACTGCCGATTCTATGAGGAA  
481 TTTGATGCCCATGGGCGTGAGGTGCCCTCCCGCTGGCATCTACAACCTGGATGACCTG  
541 AAGGCCCTGGGGCGGCGCCAGGGCTGGTGCCATACTTCCTTGCTCGATACTCAATCCTG  
601 CATGCCAATGTGGTGGTTTATAGCTACCACTACCTCCTGGACCCCAAGATTGCAGACCTG  
661 GTGTCCAAGGAACTGGCCCGCAAGGCCGTCGTGGTCTTCGACGAGGCCCAACATTGAC  
721 AACGTCTGCATCGACTCCATGAGCGTCAACCTCACCCGCCGGACCCTTGACCGGTGCCAG  
781 GGCAACCTGGAGACCCTGCAGAAGACGGTGCTCAGGATCAAAGAGACAGACGAGCAGCGC  
841 CTGCGGGACGAGTACCGGCGTCTGGTGGAGGGGCTGCGGGAGGCCAGCGCCGCCCGGGAG  
901 ACGGACGCCACCTGGCCAACCCCGTGCTGCCCGACGAAGTGCTGCAGGAGGCAGTGCCT  
961 GGCTCCATCCGCACGGCCGAGCATTTCTGGGCTTCCTGAGGCGGCTGCTGGAGTACGTG  
1021 AAGTGGCGGCTGCGTGTGCAGCATGTGGTGCAGGAGAGCCCGCCCGCCTTCCTGAGCGGC  
1081 CTGGCCCAGCGCGTGTGCATCCAGCGCAAGCCCTCAGATTCTGTGCTGAACGCCTCCGG  
1141 TCCCTGCTGCATACTCTGGAGATCACCGACCTTGCTGACTTCTCCCCGCTCACCCCTCCTT  
1201 GCTAACTTTGCCACCCTTGTCAGCACCTACGCCAAAGGCTTCACCATCATCATCGAGCCC  
1261 TTTGACGACAGAACCCCGACCATTGCCAACCCCATCCTGCACTTCAGCTGCATGGACGCC  
1321 TCGCTGGCCATCAAACCCGTATTTGAGCGTTTCCAGTCTGTATCATCACATCTGGGACA  
1381 CTGTCCCCGCTGGACATCTACCCCAAGATCCTGGACTTCACCCCGCTACCATGGCAACC  
1441 TTCACCATGACGCTGGCACGGGTCTGCCTCTGCCCTATGATCATCGGCCGTGGCAATGAC  
1501 CAGGTGGCCATCAGCTCCAAATTTGAGACCCGGGAGGATATTGCTGTGATCCGGAACAT  
1561 GGGAACCTCCTGCTGGAGATGTCCGCTGTGGTCCCTGATGGCATCGTGGCCTTCTTCACC  
1621 AGCTACCAGTACATGGAGAGCACCGTGCCCTCCTGGTATGAGCAGGGGATCCTTGAGAAC  
1681 ATCCAGAGGAACAAGCTGCTCTTTATTGAGACCCAGGATGGTGCCGAAACCAGTGTGCC  
1741 CTGGAGAAGTACCAGGAGGCCTGCGAGAATGGCCGCGGGGCCATCCTGCTGTGAGTGGCC  
1801 CGGGGCAAAGTGTCCGAGGGAATCGACTTTGTGCACCACTACGGGCGGGCCGTGATCATG  
1861 TTTGGCGTCCCCTACGTCTACACACAGAGCCGCAATTCTCAAGGCGCGGCTGGAATACCTG  
1921 CGGGACCAGTTCCAGATTCTGTGAGAATGACTTTCTTACCTTCGATGCCATGCGCCACGCG  
1981 GCCCAGTGTGTGGGTGCGGCCATCAGGGGCAAGACGGACTACGGCCTCATGGTCTTTGCC  
2041 GACAAGCGGTTTGCCCGTGCGGACAAGCGGGGAAGCTGCCCCGCTGGATCCAGGAGCAC  
2101 CTCACAGATGCCAACCTCAACCTGACCGTGGACGAGGGTGTCCAGGTGGCCAAGTACTTC  
2161 CTGCGGCAGATGGCACAGCCCTTCACCGGGAGGATCAGCTGGGCCTGTCCCTGCTCAGC  
2221 CTGGAGCAGCTAGAATCAGAGGAGACGCTGAAGAGGATAGAGCAGATTGCTCAGCAGCTC  
2281 TGAGTGGGGCGGGTGGGGCCATAAACGGTTCCTGGTGA

**Figure 4D**

MKLNVDGLLVYFPYDYIYPEQFSYMRELKRTLDAKGHGVLEMPSGTGKTVSLLALIMAYQRAYPLE  
VTKLIYCSRTVPEIEKVIEELRKLLNFYEKQEGEKL PFLGLALSSRKNL CIHPEVTPLRFGKDVDG  
KCHSLTASYVRAQYQHDTSLPHCRFYEEFDAHGREVPLPAGIYNLDDLKALGRRQGWCPYFLARYS  
ILHANVVVYSYHYLLDPKIADLVSKELARKAVVVFDEAHNIDNVCIDSMSVNLTRRTLDR CQGNLE  
TLQKTVLRIKETDEQRLRDEYRRLVEGLREASAARETDAHLANPVL PDEVLQEAVPGSIRTAEHFL  
GFLRRLLEYVKWRLRVQHVVQESPPAFLSGLAQRVCIQRKPLRFCAERLRSLLHTLEITDLADFSP  
LTL LANFATLVSTYAKGFTIIIEPFDDRTPTIANPILHFSCMDASLA IKPVFERFQSVIITSGTLS  
PLDIYPKILDFHPVTMATFTMTLARVCLCPMIIGRGNDQVAISSKFETREDIAVIRNYGNLLEMS  
AVVPDGIVAFFTSYQYMESTVASWYEQGILENIQRNKLLFIETQDGAETSVALEKYQEACENGRGA  
ILLSVARGKVSEGIDFVHHYGRAVIMFGVPYVYTQSRILKARLEYLRDQFQIRENDFLTFDAMRHA  
AQCVGRAIRGKTDYGLMVFADKRFARGDKRGKLP RWIQEHLTDANLNLTVDEGVQVAKYFLRQMAQ  
PFHREDQLGLSLLSLEQLESEETLKRIEQIAQQL

Figure 5A

CAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATAC  
ATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATTGAA  
AAAGGAAGAGTATGAGTATTCAACATTTCCGTGTGCGCCCTTATTCCTTTTTTTCGCGGCAT  
TTTGCCCTTCCTGTTTTTGTCTACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGAAGATC  
AGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGGTAAGATCCTTGAGA  
GTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCG  
CGGTATTATCCCGTATTGACGCCGGGCAAGAGCAACTCGGTGCGCCGCATACACTATTCTC  
AGAATGACTTGGTTGAGTACTCACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAG  
TAAGAGAATTATGCAGTGCTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTC  
TGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATG  
TAACTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTG  
ACACCACGATGCCTGTAGCAATGGCAACAACGTTGCGCAAACCTATTAAGTGGCGAACTAC  
TACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGGATAAAGTTGCAGGAC  
CACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGGAGCCGGTG  
AGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCG  
TAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTG  
AGATAGGTGCCTCACTGATTAAGCATTGGTAAGTGTGACACCAAGTTTACTCATATATAC  
TTTAGATTGATTTAAAACCTTCATTTTTTAATTTAAAGGATCTAGGTGAAGATCCTTTTTG  
ATAATCTCATGACCAAATCCCTTAACGTGAGTTTTCGTTCCACTGAGCGTCAGACCCCG  
TAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTCTGCGCGTAATCTGCTGCTTGC  
AAACAAAAAAACCACCGCTACCAGCGGTGGTTTTGTTTGGCGGATCAAGAGCTACCAACTC  
TTTTTCCGAAGGTAAGTGGCTTCAGCAGAGCGCAGATACCAAATACTGTCCTTCTAGTGT  
AGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTACATACCTCGCTCTGC  
TAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACT  
CAAGACGATAGTTACCGGATAAGGCGCAGCGGTGCGGCTGAACGGGGGGTTCTGTCACAC  
AGCCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAG  
AAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCG  
GAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCTG  
TCGGGTTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGGA  
GCCTATGGAAAAACGCCAGCAACGCGGCCTTTTTACGGTTCCTGGCCTTTTGCTGGCCTT  
TTGCTCACATGTTCTTTCTGCGTTATCCCTGATTCTGTGGATAACCGTATTACCGCCT  
TTGAGTGAGCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAGTCAGTGAGCG  
AGGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCCCGCGCTTGGCCGATTCAAT  
AATGCAGCTGGCACGACAGGTTTCCCGACTGGAAAGCGGGCAGTGAGCGCAACGCAATTA  
ATGTGAGTTAGCTCACTCATTAGGCACCCAGGCTTTACACTTTATGCTTCCGGCTCGTA  
TGTTGTGTGGAATTGTGAGCGGATAACAATTTACACAGGAAACAGCTATGACCATGATT  
ACGCCAAGCGCGCAATTAACCTCACTAAAGGGAACAAAAGCTGGAGCTGCAAGCTTAAT  
GTAGTCTTATGCAATACTCTTGTAGTCTTGCAACATGGTAACGATGAGTTAGCAACATGC  
CTTACAAGGAGAGAAAAAGCACCGTGATGCCGATTGGTGGAAGTAAGGTGGTACGATCG  
TGCCTTATTAGGAAGGCAACAGACGGGTCTGACATGGATTGGACGAACCACTGAATTGCC  
GCATTGCAGAGATATTGTATTTAAGTGCCTAGCTCGATACAATAAACGGGTCTCTCTGGT  
TAGACCAGATCTGAGCCTGGGAGCTCTCTGGCTAACTAGGGAACCCACTGCTTAAGCCTC  
AATAAAGCTTGCCTTGAGTGCTTCAAGTAGTGTGTGCCCGTCTGTTGTGTGACTCTGGTA  
ACTAGAGATCCCTCAGACCCTTTTAGTCAGTGTGGAAAATCTCTAGCAGTGGCGCCCGAA  
CAGGGACCTGAAAGCGAAAGGGAAACCAGAGCTCTCTCGACGCAGGACTCGGCTTGCTGA  
AGCGCGCACGGCAAGAGGCGAGGGGCGGCGACTGGTGAGTACGCCAAAAATTTTGACTAG  
CGGAGGCTAGAAGGAGAGAGATGGGTGCGAGAGCGTCAGTATTAAGCGGGGGGAGAATTAG  
ATCGCGATGGGAAAAAATTCGGTTAAGGCCAGGGGGGAAAGAAAAAATATAAATTAAACA  
TATAGTATGGGCAAGCAGGGAGCTAGAACGATTGCGAGTTAATCCTGGCCTGTTAGAAAC  
ATCAGAAGGCTGTAGACAAATACTGGGACAGCTACAACCATCCCTTCAGACAGGATCAGA

Figure 5B

AGAACTTAGATCATTATATAATACAGTAGCAACCCTCTATTGTGTGCATCAAAGGATAGA  
GATAAAAGACACCAAGGAAGCTTTAGACAAGATAGAGGAAGAGCAAAACAAAAGTAAGAC  
CACCGCACAGCAAGCGGCCGCTGATCTTCAGACCTGGAGCGCTCGAGGCGACTTACCTCT  
CTAGAGTCGGTGTCTTCTATGGAGGTCAAAACAGCGTGGATGGCGTCTCCAGGCGATCTG  
ACGGTTCACTAAACGAGCTCTGCTTATATAGACCTCCACCGTACACGCCTACCGCCCAT  
TTGCGTCAATGGGGCGGAGTTGTTACGACATTTTGAAAGTCCCGTTGATTTTGGTGCCA  
AAACAAACTCCCATTGACGTCAATGGGGTGGAGACTTGGAATCCCCGTGAGTCAAACCG  
CTATCCACGCCCATTTGATGTACTGCCAAAACCGCATCACCATGGTAATAGCGATGACTAA  
TACGTAGATGTACTGCCAAGTAGGAAAGTCCCATAGGTCATGTACTGGGCATAATGCCA  
GGCGGGCCATTTACCGTCATTGACGTCAATAGGGGGCGTACTTGGCATATGATACACTTG  
ATGTACTGCCAAGTGGGCAGTTTACCGTAAATACTCCACCCATTGACGTCAATGGAAAGT  
CCCTATTGGCGTTACTATGGGAACATACGTCATTATTGACGTCAATGGGCGGGGGTCGTT  
GGGCGGTACGCCAGGCGGGCCATTTACCGTAAGTTATGTAACGCGGAACCTCCAAGCTTA  
TCGAGGAGGAGATATGAGGGACAATTGGAGAAGTGAATTATATAAATATAAAGTAGTAAA  
AATTGAACCATTAGGAGTAGCACCCACCAAGGCAAAGAGAAGAGTGGTGCAGAGAGAAAA  
AAGAGCAGTGGGAATAGGAGCTTTGTTCCCTTGGGTTCTTGGGAGCAGCAGGAAGCACTAT  
GGGCGCAGCCTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCA  
GCAGCAGAACAAATTTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCAACTCACAGT  
CTGGGGCATCAAGCAGCTCCAGGCAAGAATCCTGGCTGTGGAAAGATACCTAAAGGATCA  
ACAGCTCCTGGGGATTTGGGGTTGCTCTGGAAAACCTCATTTGCACCACTGCTGTGCCTTG  
GAATGCTAGTTGGAGTAATAAATCTCTGGAACAGATTGGAATCACACGACCTGGATGGAG  
TGGGACAGAGAAATTAACAATTACACAAGCTTAATACACTCCTTAATTGAAGAATCGCAA  
AACCAGCAAGAAAAGAATGAACAAGAATTATTGGAATTAGATAAATGGGCAAGTTTGTGG  
AATTGGTTTAAACATAACAAATTGGCTGTGGTATATAAAATTATTCATAATGATAGTAGGA  
GGCTTGGTAGGTTTAAAGAATAGTTTTTGTCTGTACTTTCTATAGTGAATAGAGTTAGGCAG  
GGATATTCACCATTTATCGTTTCAGACCCACCTCCCAACCCCGAGGGGACCCGACAGGCCC  
GAAGGAATAGAAGAAGAAGGTGGAGAGAGAGACAGAGACAGATCCATTTCGATTAGTGAAC  
GGATCTCGACGGTTAACTTTTAAAGAAAAGGGGGGATTGGGGGGTACAGTGCAGGGGAA  
AGAATAGTAGACATAATAGCAACAGACATACAACTAAAGAATTACAAAAACAAATTACA  
AAAATTCAAATTTTATCGCATGTTCTTTCCTGCGTTATCCCCTGATTCTGTGGATAACC  
GTATTACCGCCATGCATTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGC  
CCATATATGGAGTTCGCGTTACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCC  
AACGACCCCCGCCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGG  
ACTTTCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGTACAT  
CAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCC  
TGGCATTATGCCAGTACATGACCTTATGGGACTTTCCTACTTGGCAGTACATCTACGTA  
TTAGTCATCGCTATTACCATGGTGATGCGGTTTTTGGCAGTACATCAATGGGCGTGGATAG  
CGGTTTGACTCACGGGGATTTCCAAGTCTCACCCCATTTGACGTCAATGGGAGTTTGT  
TGGCACCAAAATCAACGGGACTTTCCAAAATGTCGTAACAACTCCGCCCCATTGACGCAA  
ATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTGGTTTAGTGAACCGT  
CAGATCCGCTAGCGCTACCGGACTCAGATCTCGAGCTCAAGCTTCGAATTCTGCAGTCGA  
CGGTACCGCGGGGCCGGGATCCACCGGTGCGCCACCATGGCCTCCTCCGAGAACGTCATCA  
CCGAGTTCATGCGCTTCAAGGTGCGCATGGAGGGCACCGTGAACGGCCACGAGTTCGAGA  
TCGAGGGCGAGGGCGAGGGCCGCCCTACGAGGGGCCACAACACCGTGAAGCTGAAGGTGA  
CCAAGGGCGGGCCCCCTGCCCTTGCCTGGGACATCCTGTCCCCCAGTTCCAGTACGGCT  
CCAAGGTGTACGTGAAGCACCCCGCCGACATCCCCGACTACAAGAAGCTGTCCTTCCCCG  
AGGGCTTCAAGTGGGAGCGCGTGATGAACTTCGAGGACGGCGGGCGTGGCGACCGTGACCC  
AGGACTCCTCCCTGCAGGACGGCTGCTTCATCTACAAGGTGAAGTTCATCGGCGTGAAGT  
TCCCCTCCGACGGCCCCGTGATGCAGAAGAAGACCATGGGCTGGGAGGCCTCCACCGAGC  
GCCTGTACCCCCGCGACGGCGTGCTGAAGGGCGAGACCCACAAGGCCCTGAAGCTGAAGG

Figure 5C

ACGGCGGCCACTACCTGGTGGAGTTCAAGTCCATCTACATGGCCAAGAAGCCCGTGCAGC  
TGCCCGGCTACTACTACGTGGACGCCAAGCTGGACATCACCTCCCACAACGAGGACTACA  
CCATCGTGGAGCAGTACGAGCGCACCGAGGGCCGCCACCACCTGTTCTGTAGCGGGGCC  
TCGACAATCAACCTCTGGATTACAAAATTTGTGAAAGATTGACTGGTATTCTTAACATATG  
TTGCTCCTTTTACGCTATGTGGATACGCTGCTTTAATGCCTTTGTATCATGCTATTGCTT  
CCCGTATGGCTTTTCAATTTCTCCTCCTTGTATAAATCCTGGTTGCTGTCTCTTTATGAGG  
AGTTGTGGCCCGTTGTGAGGCAACGTGGCGTGGTGTGCACTGTGTTTGCTGACGCAACCC  
CCACTGGTTGGGGCATTGCCACCACCTGTCAGCTCCTTTCCGGGACTTTCGCTTTCCCCC  
TCCCTATTGCCACGGCGGAACTCATCGCCGCCTGCCTTGCCCGCTGCTGGACAGGGGCTC  
GGCTGTTGGGCACTGACAATTCCGTGGTGTGTGCGGGGAAGCTGACGTCCTTTCCATGGC  
TGCTCGCCTGTGTTGCCACCTGGATTCTGCGCGGGACGTCCTTCTGCTACGTCCTTCGG  
CCCTCAATCCAGCGGACCTTCCTTCCCGCGGCCTGCTGCCGGCTCTGCGGCCTCTTCGGC  
GTCTTCGCCTTCGCCCTCAGACGAGTCGGATCTCCCTTTGGGCGCCTCCCCGCCTGGAA  
TTCCGCGACTCTAGATCATAATCAGCCATACCACATTTGTAGAGGTTTTACTTGCTTTAA  
AAAACCTCCCACACCTCCCCCTGAACCTGAAACATAAAATGAATGCAATTGTTGTTGTTA  
ACTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATTCACAA  
ATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGTTTGTCCAAACTCATCAATGTATCTT  
AACCAGGCGGGGAGGCGGCCCAAAGGGAGATCCGACTCGTCTGAGGGCGAAGGCGAAGAC  
GCGGAAGAGGCCGCGAGAGCCGGCAGCAGGCCGCGGGAAGGAAGGTCCGCTGGATTGAGGG  
CCGAAGGGACGTAGCAGAAGGACGTCCCGCGCAGAATCCAGGTGGCAACACAGGCGAGCA  
GCCATGGAAAGGACGTGAGCTTCCCCGACAACACCACGGAATTGTCAGTGCCCAACAGCC  
GAGCCCCGTGTCCAGCAGCGGGCAAGGCAGGCGGCGATGAGTTCCGCCGTGGCAATAGGGA  
GGGGGAAAGCGAAAGTCCCGGAAAGGAGCTGACAGGTGGTGGCAATGCCCCAACCAGTGG  
GGGTTGCGTACGCAAACACAGTGCACACCACGCCACGTTGCCTGACAACGGGGCCACAACT  
CCTCATAAAGAGACAGCAACCAGGATTTATACAAGGAGGAGAAAATGAAAGCCATACGGG  
AAGCAATAGCATGATACAAAGGCATTAAAGCAGCGTATCCACATAGCGTAAAGGAGCAA  
CATAGTTAAGAATACCAGTCAATCTTTCACAAATTTTGTAAATCCAGAGGTTGATTGTGCA  
CGCGGCCGCTTTACTTGTACAGCTCGTCCATGCCGAGAGTGATCCCGCGGCGGGTCACGA  
ACTCCAGCAGGACCATGTGATCGCGCTTCTCGTTGGGGTCTTTGCTCAGGGCGGACTGGG  
TGCTCAGGTAGTGGTTGTGCGGCAGCAGCACGGGGCCGTGCGCGATGGGGGTGTTCTGCT  
GGTAGTGGTTCGGCGAGCTGCACGCTGCCGTCTCGATGTTGTGGCGGATCTTGAAGTTCA  
CCTTGATGCCGTTCTTCTGCTTGTGCGCCATGATATAGACGTTGTGGCTGTTGTAGTTGT  
ACTCCAGCTTGTGCCCCAGGATGTTGCCGTCTCCTTGAAGTCGATGCCCTTCAGCTCGA  
TGCGGTTTACCAGGGTGTGCCCCCGAACTTCACCTCGGCGCGGGTCTTGTAGTTGCCGT  
CGTCCTTGAAGAAGATGGTGCCTCCTGGACGTAGCCTTCGGGCATGGCGGACTTGAAGA  
AGTCGTGCTGCTTCATGTGGTCCGGGTAGCGGCTGAAGCACTGCACGCCGTAGGTGAGGG  
TGGTCACGAGGGTGGGCCAGGGCACGGGCAGCTTGCCGGTGGTGCAGATGAACTTCAGGG  
TCAGCTTGCCGTAGGTGGCATCGCCCTCGCCCTCGCCGGACACGCTGAACTTGTGGCCGT  
TTACGTGCGCGTCCAGCTCGACCAGGATGGGCACCACCCCGGTGAACAGCTCCTCGCCCT  
TGCTCACCATGGTGGCGACCGGTGGATCCTGAAGAAAAGGGAGAATTCGAATTCGAGCTC  
GGTACCTTTAAGACCAATGACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTAAAGA  
AAAGGGGGGACTGGAAGGGCTAATTCCTCCCAACGAAGACAAGATCTGCTTTTTGCTTG  
TACTGGGTCTCTCTGGTTAGACCAGATCTGAGCCTGGGAGCTCTCTGGCTAACTAGGGAA  
CCCCTGCTTAAGCCTCAATAAAGCTTGCCTTGAGTGCTTCAAGTAGTGTGTGCCCGTCT  
GTTGTGTGACTCTGGTAAGTAGAGATCCCTCAGACCCTTTTAGTCAGTGTGGAAAATCTC  
TAGCAGTAGTAGTTCATGTCATCTTATTATTCAGTATTTATAACTTGCAAAGAAATGAAT  
ATCAGAGAGTGAGAGGAACCTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAG  
CATCACAAATTTACAAATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGTTTGTCCAA  
ACTCATCAATGTATCTTATCATGTCTGGCTCTAGCTATCCCGCCCCCTAACTCCGCCAGT  
TCCGCCCATCTCCGCCCATGGCTGACTAATTTTTTTTTTATTATGCAGAGGCCGAGGCC

**Figure 5D**

GCCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTT  
TGCGTCGAGACGTACCCAATTCGCCCTATAGTGAGTCGTATTACGCGCGCTCACTGGCCG  
TCGTTTTACAACGTCGTGACTGGGAAAACCCTGGCGTTACCCAACCTAATCGCCTTGCAG  
CACATCCCCCTTTCGCCAGCTGGCGTAATAGCGAAGAGGCCCGCACCGATCGCCCTTCCC  
AACAGTTGCGCAGCCTGAATGGCGAATGGCGCGACGCGCCCTGTAGCGGCGCATTAAGCG  
CGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCCG  
CTCCTTTCGCTTTCCTTCCCTTCCTTTCCTCGCCACGTTTCGCCGGCTTTCCTCCGTCAAGCTC  
TAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCAAAA  
AACTTGATTAGGGTGATGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTTCGCC  
CTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAAACTGGAACAACAC  
TCAACCCTATCTCGGTCTATTCTTTTGATTTATAAGGGATTTTGCCGATTTTCGGCCTATT  
GGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGCGAATTTTAACAAAATATTAAAGT  
TTACAATTTCC

Figure 6A

CCGGTCGCCACCATGGCCTCCTCCGAGAACGTCATCACCGAGTTCATGCGCTTCAAGGTGCGCA  
TGGAGGGCACCCTGAACGGCCACGAGTTCGAGATCGAGGGCGAGGGCGAGGGCCGCCCTACGA  
GGGCCACAACACCGTGAAGCTGAAGGTGACCAAGGGCGGCCCCCTGCCCTTCGCTGGGACATC  
CTGTCCCCCAGTTCAGTACGGCTCCAAGGTGTACGTGAAGCACCCCGCCGACATCCCCGACT  
ACAAGAAGCTGTCCTTCCCCGAGGGCTTCAAGTGGGAGCGCGTGATGAACCTCGAGGACGGCGG  
CGTGGCGACCGTGACCCAGGACTCCTCCCTGCAGGACGGCTGCTTCATCTACAAGGTGAAGTTC  
ATCGGCGTGAACCTTCCCCTCCGACGGCCCCGTGATGCAGAAGAAGACCATGGGCTGGGAGGCCT  
CCACCGAGCGCCTGTACCCCCGCGACGGCGTGCTGAAGGGCGAGACCCACAAGGCCCTGAAGCT  
GAAGGACGGCGGCCACTACCTGGTGGAGTTCAGTCCATCTACATGGCCAAGAAGCCCGTGAG  
CTGCCCGGCTACTACTACGTGGACGCCAAGCTGGACATCACCTCCACAACGAGGACTACACCA  
TCGTGGAGCAGTACGAGCGCACCGAGGGCCGCCACCACCTGTTCTGTAGCGGCCGCGACTCTA  
GATCATAATCAGCCATAACACATTTGTAGAGGTTTTACTTGCTTTAAAAAACCTCCCACACCTC  
CCCCTGAACCTGAAACATAAAATGAATGCAATTGTTGTTGTTAACTTGTTTTATTGCAGCTTATA  
ATGGTTACAAATAAAGCAATAGCATCACAAATTTACAAATAAAGCATTTTTTTTCACTGCATTC  
TAGTTGGATCCTTTATTTGTGAAATTTGTGATGCTATTGCTTTATTTGTAACCATTATAAGCTG  
CAATAAACAAGTTAACAACAACAATTGCATTCATTTTATGTTTCAGGTTCAAGGGGAGGTGTGG  
GAGGTTTTTTTAAAGCAAGTAAACCTCTACAAATGTGGTATGGCTGAATTCAGGGCGGGGAGGC  
GGCCCAAAGGGAGATCCGACTCGTCTGAGGGCGAAGGCGAAGACGCGGAAGAGGCCGAGAGCC  
GGCAGCAGGCCGCGGGAAGGAAGGTCCGCTGGATTGAGGGCCGAAGGGACGTAGCAGAAGGACG  
TCCCGCGCAGAATCCAGGTGGCAACACAGGCGAGCAGCCATGGAAAGGACGTCAGCTTCCCCGA  
CAACACCACGGAATTGTCAGTGCCCAACAGCCGAGCCCCTGTCCAGCAGCGGGCAAGGCAGGCG  
GCGATGAGTTCCGCCGTGGCAATAGGGAGGGGGAAAGCGAAAGTCCCGGAAAGGAGCTGACAGG  
TGGTGGCAATGCCCCAACCAGTGGGGGTTCGCTCAGCAAACACAGTGACACACCACGCCACGTTG  
CCTGACAACGGGGCCACAACCTCCTCATAAAGAGACAGCAACCAGGATTTATACAAGGAGGAGAAA  
ATGAAAGCCATACGGGAAGCAATAGCATGATACAAAGGCATTAAAGCAGCGTATCCACATAGCG  
TAAAAGGAGCAACATAGTTAAGAATACCAGTCAATCTTTCACAAATTTTGTAAATCCAGAGGTTG  
ATTGTCGACGCGGCCGCTTTACTTGTACAGCTCGTCCATGCCGAGAGTGATCCCGGCCGCGGTC  
ACGAACTCCAGCAGGACCATGTGATCGCGCTTCTCGTTGGGGTCTTTGCTCAGGGCGGACTGGG  
TGCTCAGGTAGTGGTTGTGCGGCAGCAGCACGGGGCCGTCCCGATGGGGGTGTTCTGCTGGTA  
GTGGTCGGCGAGCTGCACGCTGCCGTCTCGATGTTGTGGCGGATCTTGAAGTTCACCTTGATG  
CCGTTCTTCTGCTTGTGCGCCATGATATAGACGTTGTGGCTGTTGTAGTTGTACTCCAGCTTGT  
GCCCCAGGATGTTGCCGTCTCCTTGAAGTCGATGCCCTTCAGCTCGATGCGGTTCCACAGGGT  
GTCGCCCTCGAACTTCACCTCGGCGCGGGTCTTGTAGTTGCCGTCTGCTTGAAGAAGATGGTG  
CGCTCCTGGACGTAGCCTTCGGGCATGGCGGACTTGAAGAAGTCGTGCTGCTTCATGTGGTTCG  
GGTAGCGGCTGAAGCACTGCACGCCGTAGGTGAGGTGGTCACGAGGGTGGGCCAGGGCACGGG  
CAGCTTGCCGGTGGTGCAGATGAACCTCAGGGTCAGCTTGCCGTAGGTGGCATCGCCCTCGCCC  
TCGCCGGACACGCTGAACCTTGTGGCCGTTTACGTCGCCGTCCAGCTCGACCAGGATGGGCACCA  
CCCCGGTGAACAGCTCCTCGCCCTTGCTCACCATCTGAAGAAAAGGGAGGTACCTTTAAGACCA  
ATGACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTTAAAGAAAAGGGGGGACTGGAAGGGC  
TAATTCACCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACACAAGGCTACTT  
CCCTGATTGGCAGAATAACACACCAGGGCCAGGGATCAGATATCCACTGACCTTTGGATGGTGC  
TACAAGCTAGTACCAGTTGAGCAAGAGAAGGTAGAAGAAGCCAATGAAGGAGAGAACACCCGCT  
TGTTACACCCTGTGAGCCTGCATGGGATGGATGACCCGGAGAGAGAAGTATTAGAGTGGAGGTT  
TGACAGCCGCCTAGCATTTTCATCACATGGCCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGC  
TGACATCGAGCTTGCTACAAGGGACTTTCCGCTGGGGACTTTCCAGGGAGGCGTGCCCTGGGCG  
GGACTGGGGAGTGGCGAGCCCTCAGATGCTGCATATAAGCAGCTGCTTTTTTGCTTGTACTGGGT  
CTCTCTGGTTAGACCAGATCTGAGCCTGGGAGCTCTCTGGCTAACTAGGGAACCCACTGCTTAA  
GCCTCAATAAAGCTTGCCCTTGAGTGCTTCAAGTAGTGTGTGCCCGTCTGTTGTGTGACTCTGGT  
AACTAGAGATCCCTCAGACCCTTTTAGTCAGTGTGGAAAATCTCTAGCAGTAGTAGTTCATGTC  
ATCTTATTATTACAGTATTTATAACTTGCAAAGAAATGAATATCAGAGAGTGAGAGGCCTTGACA



**Figure 6B**

TTATAATAGATTTAGCAGGAATTGAACTAGGAGTGGAGCACACAGGCAAAGCTGCAGAAGTACT  
TGGAAGAAGCCACCAGAGATACTCACGATTCTGCACATACCTGGCTAATCCCAGATCCTAAGGA  
TTACATTAAGTTTACTAACATTTATATAATGATTTATAGTTTAAAGTATAAACTTATCTAATTT  
ACTATTCTGACAGATATTAATTAATCCTCAAATATCATAAGAGATGATTACTATTATCCCCATT  
TAACACAAGAGGAACTGAGAGGGAAAGATGTTGAAGTAATTTCCACAAATTACAGCATCCGT  
TAGTTACGACTCTATGATCTTCTGACACAAATTCCATTTACTCCTCACCTATGACTCAGTCGA  
ATATATCAAAGTTATGGACATTATGCTAAGTAACAAATTACCCTTTTATATAGTAAATACTGAG  
TAGATTGAGAGAAGAAATTGTTTGGCAAACCTGAATAGCTTCCAGAAGAAGAGAAGTGAGGATA  
AGAATAACAGTTGTCATTAACCAGTTTTTAACAAGTAACCTTGGTTAGAAAGGGATTCAAATGCAT  
AAAGCAAGGGATAAATTTTTCTGGCAACAAGACTATACAATATAACCTTAAATATGACTTCAA  
TAATTGTTGGAACCTTGATAAACTAATTAATATTATTGAAGATTATCAATATTATAAATGTAA  
TTTACTTTTAAAAAGGGAACATAGAAATGTGTATCATTAGAGTAGAAAACAATCCTTATTATCA  
CAATTTGTCAAACAAGTTTGTATTAAACAAGTAGAATACTGCATTCAATTAAGTTGACTGC  
AGATTTTGTGTTTTGTAAATTAGAAAGAGATAACAACAATTTGAATTATTGAAAGTAACATG  
TAAATAGTTCTACATACGTTCTTTTGACATCTTGTTCATCATTGATCGAAGTTCTTTATCTTG  
GAAGAATTTGTTCCAAAGACTCTGAAATAAGGAAAACAATCTATTATATAGTCTCACACCTTTG  
TTTTACTTTTGTGATTTCAATTTAATAATGTAAATGGTTAAAATTTATTCTTCTCTGAGATCA  
TTTCACATTGCAGATAGAAAACCTGAGACTGGGGTAATTTTTTATTAAAATCTAATTTAATCTCA  
GAAACACATCTTTATTCTAACATCAATTTTTCCAGTTTGATATTATCATATAAAGTCAGCCTTC  
CTCATCTGCAGGTTCCACAACAAAAATCCAACCACTGTGGATCAAAAATATTGGGAAAAAATT  
AAAAATAGCAATACAACAATAAAAAAATACAAATCAGAAAAACAGCACAGTATAACAACCTTTAT  
TTAGCATTTACAATCTATTAGGTATTATAAGTAATCTAGAATTAATTCCGTGTATTCTATAGTG  
TCACCTAAATCGTATGTGTATGATACATAAGGTTATGTATTAATTGTAGCCGCGTTCTAACGAC  
AATATGTACAAGCCTAATTGTGTAGCATCTGGCTTACTGAAGCAGACCCTATCATCTCTCTCGT  
AAACTGCCGTCAGAGTCGGTTTTGTTTGGACGAACCTTCTGAGTTTCTGGTAACGCCGTTCCGCA  
CCCCGGAAATGGTCAGCGAACCAATCAGCAGGGTCATCGCTAGCCAGATCCTCTACGCCGGACG  
CATCGTGGCCGGCATCACCGGCGCCACAGGTGCGGTTGCTGGCGCCTATATCGCCGACATCACC  
GATGGGGAAGATCGGGCTCGCCACTTCGGGCTCATGAGCGCTTGTTTCGGCGTGGGTATGGTGG  
CAGGCCCCGTGGCCGGGGGACTGTTGGGCGCCATCTCCTTGATGCACCATTCTTGCGGCGGC  
GGTGCTCAACGGCCTCAACCTACTACTGGGCTGCTTCTAATGCAGGAGTCGCATAAGGGAGAG  
CGTCGATATGGTGCACTCTCAGTACAATCTGCTCTGATGCCGCATAGTTAAGCCAGCCCCGACA  
CCCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCCGGCATCCGCTTACAGACAA  
GCTGTGACCGTCTCCGGGAGCTGCATGTGTGAGAGGTTTTACCGTCATCACCGAAACGCGCGA  
GACGAAAGGGCCTCGTGATACGCCATTTTTTATAGGTTAATGTCATGATAATAATGGTTTCTTA  
GACGTGAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTTATTTTTCTAAATA  
CATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAA  
GGAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCTTTTTTTCGGGCATTTTGCCT  
TCCTGTTTTTGTCTACCCAGAAACGCTGGTGAAAGTAAAGATGCTGAAGATCAGTTGGGTGCA  
CGAGTGGGTTACATCGAACTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAG  
AACGTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGA  
CGCCGGGCAAGAGCAACTCGGTGCGCGCATACACTATTCTCAGAATGACTTGTTGAGTACTCA  
CCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAA  
CCATGAGTGATAACACTGCGGCCAATTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAAC  
CGCTTTTTTGCACAACATGGGGGATCATGTAACCTGCCTTGATCGTTGGGAACCGGAGCTGAAT  
GAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTAGCAATGGCAACAACGTTGCGCA  
AACTATTAACCTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGC  
GGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTTATTGCTGATAAA  
TCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCT

Figure 6C

CCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGAT  
CGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTCAGACCAAGTTTACTCATATATA  
CTTTAGATTGATTTAAACTTCATTTTTTAATTTAAAGGATCTAGGTGAAGATCCTTTTTTGATA  
ATCTCATGACCAAATCCCTTAACGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAA  
GATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTGCTGCTTGCAAACAAAAAA  
CCACCGCTACCAGCGGTGGTTTGTGTGCGGATCAAGAGCTACCAACTCTTTTTCCGAAGGTAA  
CTGGCTTCAGCAGAGCGCAGATACCAAATACTGTCTTCTAGTGTAGCCGTAGTTAGGCCACCA  
CTTCAAGAACTCTGTAGCACCGCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCT  
GCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGC  
AGCGGTCGGGCTGAACGGGGGGTTCGTGCACACAGCCAGCTTGGAGCGAACGACCTACACCGA  
ACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGAC  
AGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACG  
CCTGGTATCTTTATAGTCTGTGCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTTGTGATG  
CTCGTCAGGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGCGGCCTTTTTTACGGTTCCTGGCC  
TTTTGCTGGCCTTTTGCTCACATGTTCTTTCCTGCGTTATCCCCTGATTCTGTGGATAACCGTA  
TTACCGCCTTTGAGTGAGCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAGCGAGTCAGT  
GAGCGAGGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCCCGCGCGTTGGCCGATTTCAT  
TAATGCAGCTGTGGAATGTGTGTGTCAGTTAGGGTGTGGAAAGTCCCCAGGCTCCCCAGCAGGCAG  
AAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCAGGTGTGGAAAGTCCCCAGGCTCCCCA  
GCAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTC  
CGCCCATCCCGCCCCCTAACTCCGCCCAGTTCGCCCCATTCTCCGCCCCATGGCTGACTAATTTT  
TTTTATTTATGCAGAGGCCGAGGCCGCTCGGCCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGC  
TTTTTTGGAGGCCCTAGGCTTTTGCAAAAAGCTTGGACACAAGACAGGCTTGCGAGATATGTTTG  
AGAATACCACTTTATCCCGCGTCAGGGAGAGGCAGTGCGTAAAAAGACGCGGACTCATGTGAAA  
TACTGGTTTTTTAGTGCGCCAGATCTCTATAATCTCGCGCAACCTATTTTCCCCTCGAACACTTT  
TTAAGCCGTAGATAAACAGGCTGGGACACTTCACATGAGCGAAAAATACATCGTCACCTGGGAC  
ATGTTGCAGATCCATGCACGTAACTCGCAAGCCGACTGATGCCTTCTGAACAATGGAAAGGCA  
TTATTGCCGTAAGCCGTGGCGGTCTGTACCGGGTGCCTTACTGGCGCGTGAACCTGGGTATTTCGT  
CATGTCGATACCGTTTGTATTTCCAGCTACGATCACGACAACCAGCGCGAGCTTAAAGTGCTGA  
AACGCGCAGAAGGCGATGGCGAAGGCTTCATCGTTATTGATGACCTGGTGGATACCGGTGGTAC  
TGCGGTTGCGATTCTGTGAAATGTATCCAAAAGCGCACTTTGTACCATCTTCGCAAAACCGGCT  
GGTCGTCCGCTGGTTGATGACTATGTTGTTGATATCCCGCAAGATACCTGGATTGAACAGCCGT  
GGGATATGGGCGTCGTATTCGTCCCGCCAATCTCCGGTCGCTAATCTTTTCAACGCCTGGCACT  
GCCGGGCGTTGTTCTTTTTAACTTCAGGCGGGTTACAATAGTTTCCAGTAAGTATTCTGGAGGC  
TGCATCCATGACACAGGCAAACCTGAGCGAAACCCTGTTCAAACCCCGCTTTAAACATCCTGAA  
ACCTCGACGCTAGTCCGCGCTTTAATCACGGCGCACAAACCGCTGTGCAGTCGGCCCTTGATG  
GTAAAACCATCCCTCACTGGTATCGCATGATTAACCGTCTGATGTGGATCTGGCGCGGCATTGA  
CCCACGCGAAATCCTCGACGTCCAGGCACGTATTGTGATGAGCGATGCCGAACGTACCGACGAT  
GATTTATACGATACGGTGATTGGCTACCGTGGCGGCAACTGGATTTATGAGTGGGCCCCGGATC  
TTTGTGAAGGAACCTTACTTCTGTGGTGTGACATAATTGGACAAACTACCTACAGAGATTTAAA  
GCTCTAAGGTAAATATAAAATTTTTAAGTGTATAATGTGTTAAACTACTGATTCTAATTGTTTG  
TGTATTTTAGATTCCAACCTATGGAAGTGAATGGGAGCAGTGGTGAATGCCTTTAATGAG  
GAAAACCTGTTTTGCTCAGAAGAAATGCCATCTAGTGATGATGAGGCTACTGCTGACTCTCAAC  
ATTCTACTCCTCCAAAAAAGAAGAGAAAGGTAGAAGACCCCAAGGACTTTCCTTCAGAATTGCT  
AAGTTTTTTGAGTCATGCTGTGTTTAGTAATAGAACTCTTGCTTGCTTTGCTATTTACACCACA  
AAGGAAAAAGCTGCACTGCTATAAGAAAATTATGGAATAATTCTGTAACCTTTATAAGTA  
GGCATAACAGTTATAATCATAACATACTGTTTTTTCTTACTCCACACAGGCATAGAGTGTCTGC  
TATTAATAACTATGCTCAAAAATTGTGTACCTTTAGCTTTTTAATTTGTAAAGGGGTTAATAAG  
GAATATTTGATGTATAGTGCCTTGACTAGAGATCATAATCAGCCATACCACATTTGTAGAGGTT  
TTACTTGCTTTAAAAAACCTCCCACACCTCCCCCTGAACCTGAAACATAAAATGAATGCAATTG

**Figure 6D**

TTGTTGTTAACTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATTT  
CACAAATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGTTTGTCCAACTCATCAATGTATCT  
TATCATGTCTGGATCAACTGGATAACTCAAGCTAACCAAAATCATCCCAAACCTCCCACCCCAT  
ACCCTATTACCACTGCCAATTACCTGTGGTTTCATTTACTCTAAACCTGTGATTCCTCTGAATT  
ATTTTCATTTTAAAGAAATTGTATTTGTTAAATATGTACTACAACTTAGTAGT

## SEQUENCE LISTING

<110> Fishel, Richard A.  
Yoder, Kristine E.

<120> Methods Of Identifying Compounds That Modulate A DNA Repair  
Pathway And/or Retroviral Infectivity, The Compounds, And Uses  
Thereof

<130> SCHN0003

<160> 6

<170> PatentIn version 3.2

<210> 1  
<211> 2751  
<212> DNA  
<213> Homo sapiens

<400> 1  
 gggagcttcc ggattgagcc ggaagtcccc ccagagcgga tgccgcggcg ggcctgtggg 60  
 agcgggggtca tcttctctct gctgctgtag ctgccatggg caaaagagac cgagcggacc 120  
 gcgacaagaa gaaatccagg aagcggcact atgaggatga agaggatgat gaagaggacg 180  
 ccccggggaa cgaccctcag gaagcggttc cctcggcggc ggggaagcag gtggatgagt 240  
 caggcaccaa agtggatgaa tatggagcca aggactacag gctgcaaatg ccgctgaagg 300  
 acgaccacac ctccaggccc ctctgggtgg ctcccgatgg ccatatcttc ttggaagcct 360  
 tctctccagt ttacaaatat gcccaagact tcttggtggc tattgcagag ccagtgtgcc 420  
 gaccaaccca tgtgcatgag tacaaactaa ctgcctactc cttgtatgca gctgtcagcg 480  
 ttgggtgca aaccagtgc atcaccgagt acctcaggaa gctcagcaag actggagtcc 540  
 ctgatggaat tatgcagttt attaagttgt gtactgtcag ctatggaaaa gtcaagctgg 600  
 tcttgaagca caacagatac ttcgttgaaa gttgccaccc tgatgtaatc cagcatcttc 660  
 tccaggaccc cgtgatccga gaatgccgct taagaaactc tgaaggggag gccactgagc 720  
 tcatcacaga gactttcaca agcaaactct ccatctctaa gactgctgaa agcagtgggtg 780  
 ggccctccac ttcccagtg acagatccac agggtaaatac tgacatcccc atggacctgt 840  
 ttgacttcta tgagcaaatg gacaaggatg aagaagaaga agaagagaca cagacagtgt 900  
 cttttgaagt caagcaggaa atgattgagg aactccagaa acgttgcatc cacctggagt 960  
 accctctgtt ggcagaatat gacttccgga atgattctgt caaccctgat atcaacattg 1020  
 acctaaagcc cacagctgtc ctccagaccct atcaggagaa gagcttgca aagatgtttg 1080  
 gaaacgggcg tgcacgttcg ggggtcattg ttcttccctg cgggtgctgga aagtccctgg 1140  
 ttggtgtgac tgctgcatgc actgtcagaa aacgctgtct ggtgctgggc aactcagctg 1200  
 tttctgtgga gcagtggaaa gccagttca agatgtggtc caccattgac gacagccaga 1260  
 tctgccggtt cacctccgat gccaaaggaca agccatcgg ctgctccgtt gccattagca 1320  
 cctactccat gctgggccac accaccaaaa ggtcctggga ggccgagcga gtcattggagt 1380  
 ggctcaagac ccaggagtgg ggcctcatga tctggatga agtgcacacc ataccagcca 1440  
 agatgttccg aagggtgctc accatcgtgc aggccactg taagctgggt ttgactgcga 1500

```

ccctcggtccg cgaagatgac aaaattgtgg atttaaattt tctgattggg cctaagctct 1560
acgaagccaa ctggatggag ctgcagaata atggctacat cgccaaagtc cagtgtgctg 1620
aggtctggtg ccctatgtct cctgaatttt accgggaata tgtggcaatc aaaaccaaga 1680
aacgaatctt gctgtacacc atgaacccca acaaatttag agcttgccag tttctgatca 1740
agtttcatga aaggaggaat gacaagatta ttgtctttgc tgacaatgtg tttgccctaa 1800
aggaatatgc cattcgactg aacaaaccct atatctacgg acctacgtct caggggggaaa 1860
ggatgcaaat tctccagaat ttcaagcaca accccaaaat taacaccatc ttcatatcca 1920
aggtaggtga cacttcgttt gatctgccgg aagcaaatgt cctcattcag atctcatccc 1980
atggtggctc caggcgctcag gaagcccaaa ggctagggcg ggtgcttcga gctaaaaaag 2040
ggatgggttg agaagagtac aatgcctttt tctactcact ggtatcccag gacacacagg 2100
aaatggctta ctcaaccaag cggcagagat tcttggtaga tcaaggttat agcttcaagg 2160
tgatcacgaa actcgctggc atggaggagg aagacttggc gttttcgaca aaagaagagc 2220
aacagcagct cttacagaaa gtcctggcag ccactgacct ggatgccgag gaggaggtgg 2280
tggtcgggga atttggtctc agatccagcc aggcattctc gcgctttggc accatgagtt 2340
ctatgtctgg ggccgacgac actgtgtaca tggagtacca ctcatcgcg agcaaggcgc 2400
ccagcaaaca tgtacacccg ctcttcaagc gctttaggaa atgatgctta ggcagggtac 2460
ttcgttcaag accggcgctt ggcacccttg ttggaaaggg attttcagca taacattttc 2520
cttcacctc tttgaccttc cctccagcgt tggccaaatt gtgctgagga agatgcatca 2580
agggcttggc tgtgccttca taggtcatct agggttttat aaaggaggag gagacaatat 2640
tttttcaaac tttttgggga gtgggggtcat ttctgtatat aaaaaatgtt aatatttaag 2700
gtgtatttat gttaccgttc tgaataaaca gaatggacca ttgaaccagt a 2751

```

<210> 2  
 <211> 782  
 <212> PRT  
 <213> Homo sapiens

<400> 2

Met Gly Lys Arg Asp Arg Ala Asp Arg Asp Lys Lys Lys Ser Arg Lys  
 1 5 10 15

Arg His Tyr Glu Asp Glu Glu Asp Asp Glu Glu Asp Ala Pro Gly Asn  
 20 25 30

Asp Pro Gln Glu Ala Val Pro Ser Ala Ala Gly Lys Gln Val Asp Glu  
 35 40 45

Ser Gly Thr Lys Val Asp Glu Tyr Gly Ala Lys Asp Tyr Arg Leu Gln  
 50 55 60

Met Pro Leu Lys Asp Asp His Thr Ser Arg Pro Leu Trp Val Ala Pro  
 65 70 75 80

Asp Gly His Ile Phe Leu Glu Ala Phe Ser Pro Val Tyr Lys Tyr Ala

85

90

95

Gln Asp Phe Leu Val Ala Ile Ala Glu Pro Val Cys Arg Pro Thr His  
 100 105 110  
 Val His Glu Tyr Lys Leu Thr Ala Tyr Ser Leu Tyr Ala Ala Val Ser  
 115 120 125  
 Val Gly Leu Gln Thr Ser Asp Ile Thr Glu Tyr Leu Arg Lys Leu Ser  
 130 135 140  
 Lys Thr Gly Val Pro Asp Gly Ile Met Gln Phe Ile Lys Leu Cys Thr  
 145 150 155 160  
 Val Ser Tyr Gly Lys Val Lys Leu Val Leu Lys His Asn Arg Tyr Phe  
 165 170 175  
 Val Glu Ser Cys His Pro Asp Val Ile Gln His Leu Leu Gln Asp Pro  
 180 185 190  
 Val Ile Arg Glu Cys Arg Leu Arg Asn Ser Glu Gly Glu Ala Thr Glu  
 195 200 205  
 Leu Ile Thr Glu Thr Phe Thr Ser Lys Ser Ala Ile Ser Lys Thr Ala  
 210 215 220  
 Glu Ser Ser Gly Gly Pro Ser Thr Ser Arg Val Thr Asp Pro Gln Gly  
 225 230 235 240  
 Lys Ser Asp Ile Pro Met Asp Leu Phe Asp Phe Tyr Glu Gln Met Asp  
 245 250 255  
 Lys Asp Glu Glu Glu Glu Glu Thr Gln Thr Val Ser Phe Glu Val  
 260 265 270  
 Lys Gln Glu Met Ile Glu Glu Leu Gln Lys Arg Cys Ile His Leu Glu  
 275 280 285  
 Tyr Pro Leu Leu Ala Glu Tyr Asp Phe Arg Asn Asp Ser Val Asn Pro  
 290 295 300  
 Asp Ile Asn Ile Asp Leu Lys Pro Thr Ala Val Leu Arg Pro Tyr Gln  
 305 310 315 320  
 Glu Lys Ser Leu Arg Lys Met Phe Gly Asn Gly Arg Ala Arg Ser Gly  
 325 330 335  
 Val Ile Val Leu Pro Cys Gly Ala Gly Lys Ser Leu Val Gly Val Thr  
 340 345 350  
 Ala Ala Cys Thr Val Arg Lys Arg Cys Leu Val Leu Gly Asn Ser Ala  
 355 360 365

Val Ser Val Glu Gln Trp Lys Ala Gln Phe Lys Met Trp Ser Thr Ile  
 370 375 380

Asp Asp Ser Gln Ile Cys Arg Phe Thr Ser Asp Ala Lys Asp Lys Pro  
 385 390 395 400

Ile Gly Cys Ser Val Ala Ile Ser Thr Tyr Ser Met Leu Gly His Thr  
 405 410 415

Thr Lys Arg Ser Trp Glu Ala Glu Arg Val Met Glu Trp Leu Lys Thr  
 420 425 430

Gln Glu Trp Gly Leu Met Ile Leu Asp Glu Val His Thr Ile Pro Ala  
 435 440 445

Lys Met Phe Arg Arg Val Leu Thr Ile Val Gln Ala His Cys Lys Leu  
 450 455 460

Gly Leu Thr Ala Thr Leu Val Arg Glu Asp Asp Lys Ile Val Asp Leu  
 465 470 475 480

Asn Phe Leu Ile Gly Pro Lys Leu Tyr Glu Ala Asn Trp Met Glu Leu  
 485 490 495

Gln Asn Asn Gly Tyr Ile Ala Lys Val Gln Cys Ala Glu Val Trp Cys  
 500 505 510

Pro Met Ser Pro Glu Phe Tyr Arg Glu Tyr Val Ala Ile Lys Thr Lys  
 515 520 525

Lys Arg Ile Leu Leu Tyr Thr Met Asn Pro Asn Lys Phe Arg Ala Cys  
 530 535 540

Gln Phe Leu Ile Lys Phe His Glu Arg Arg Asn Asp Lys Ile Ile Val  
 545 550 555 560

Phe Ala Asp Asn Val Phe Ala Leu Lys Glu Tyr Ala Ile Arg Leu Asn  
 565 570 575

Lys Pro Tyr Ile Tyr Gly Pro Thr Ser Gln Gly Glu Arg Met Gln Ile  
 580 585 590

Leu Gln Asn Phe Lys His Asn Pro Lys Ile Asn Thr Ile Phe Ile Ser  
 595 600 605

Lys Val Gly Asp Thr Ser Phe Asp Leu Pro Glu Ala Asn Val Leu Ile  
 610 615 620

Gln Ile Ser Ser His Gly Gly Ser Arg Arg Gln Glu Ala Gln Arg Leu  
 625 630 635 640

Gly Arg Val Leu Arg Ala Lys Lys Gly Met Val Ala Glu Glu Tyr Asn  
 645 650 655

Ala Phe Phe Tyr Ser Leu Val Ser Gln Asp Thr Gln Glu Met Ala Tyr  
660 665 670

Ser Thr Lys Arg Gln Arg Phe Leu Val Asp Gln Gly Tyr Ser Phe Lys  
675 680 685

Val Ile Thr Lys Leu Ala Gly Met Glu Glu Glu Asp Leu Ala Phe Ser  
690 695 700

Thr Lys Glu Glu Gln Gln Gln Leu Leu Gln Lys Val Leu Ala Ala Thr  
705 710 715 720

Asp Leu Asp Ala Glu Glu Val Val Ala Gly Glu Phe Gly Ser Arg  
725 730 735

Ser Ser Gln Ala Ser Arg Arg Phe Gly Thr Met Ser Ser Met Ser Gly  
740 745 750

Ala Asp Asp Thr Val Tyr Met Glu Tyr His Ser Ser Arg Ser Lys Ala  
755 760 765

Pro Ser Lys His Val His Pro Leu Phe Lys Arg Phe Arg Lys  
770 775 780

<210> 3

<211> 2318

<212> DNA

<213> Homo sapiens

<400> 3

|  |     |
|--|-----|
| atgaagctca acgtggacgg gctcctggtc tacttcccgt acgactacat ctaccccgag  | 60  |
| cagttctcct acatgcggga gctcaaacgc acgctggacg ccaaggggtca tggagtcctg | 120 |
| gagatgccct caggcaccgg gaagacagta tccctgttgg ccctgatcat ggcataccag  | 180 |
| agagcatatc cgctggaggt gaccaaactc atctactgct caagaactgt gccagagatt  | 240 |
| gagaaggtga ttgaagagct tcgaaagttg ctcaacttct atgagaagca ggagggcgag  | 300 |
| aagctgccgt ttctgggact ggtcttgagc tcccgcaaaa acttgtgtat tcaccctgag  | 360 |
| gtgacacccc tgcgcttttg gaaggacgtc gatgggaaat gccacagcct cacagcctcc  | 420 |
| tatgtgcggg cgcagtacca gcatgacacc agcctgcccc actgccgatt ctatgaggaa  | 480 |
| tttgatgccc atgggctga ggtgcccctc cccgctggca tctacaacct ggatgacctg   | 540 |
| aaggccctgg ggcggcgcca gggctggtgc ccatacttcc ttgctcgata ctcaatcctg  | 600 |
| catgccaatg tgggtggtta tagctaccac tacctcctgg accccaagat tgcagacctg  | 660 |
| gtgtccaagg aactggcccg caaggccgtc gtggtcttcg acgaggccca caacattgac  | 720 |
| aacgtctgca tcgactccat gagcgtcaac ctcaccgcgc ggacccttga ccggtgccag  | 780 |
| ggcaacctgg agacctgca gaagacggtg ctcaggatca aagagacaga cgagcagcgc   | 840 |
| ctgcgggacg agtaccggcg tctggtggag gggctgcggg aggccagcgc cgcccgagg   | 900 |
| acggacgccc acctggccaa ccccgctgctg cccgacgaag tgctgcagga ggcagtgcct | 960 |



ggctccatcc gcacggccga gcatttcctg ggcttcctga ggcggctgct ggagtacgtg 1020  
 aagtggcggc tgcgtgtgca gcatgtggtg caggagagcc cggccgcctt cctgagcggc 1080  
 ctggcccagc gcgtgtgcat ccagcgcaag ccctcagat tctgtgtgta acgcctccgg 1140  
 tccctgctgc atactctgga gatcaccgac cttgctgact tctccccgct caccctcctt 1200  
 gctaactttg ccacccttgt cagcacctac gccaaaggct tcaccatcat catcgagccc 1260  
 tttgacgaca gaaccccgc catttgccaac cccatcctgc acttcagctg catggacgcc 1320  
 tcgctggcca tcaaaccgct atttgagcgt ttccagtctg tcatcatcac atctgggaca 1380  
 ctgtccccgc tggacatcta cccaagatc ctggacttcc acccgcacac catggcaacc 1440  
 ttcaccatga cgctggcagc ggtctgcctc tgccctatga tcatcggccg tggcaatgac 1500  
 caggtggcca tcagctccaa atttgagacc cgggaggata ttgctgtgat ccggaactat 1560  
 gggaacctcc tgctggagat gtccgctgtg gtccctgatg gcatcgtggc cttcttcacc 1620  
 agctaccagt acatggagag caccgtggcc tcctgggtatg agcaggggat ccttgagaac 1680  
 atccagagga acaagctgct ctttattgag acccaggatg gtgccgaaac cagtgtcgcc 1740  
 ctggagaagt accaggaggg ctgcgagaat ggccgcgggg ccatcctgct gtcagtggcc 1800  
 cggggcaaag tgtccgaggg aatcgacttt gtgcaccact acgggcgggc cgtcatcatg 1860  
 tttggcgtcc cctacgtcta cacacagagc cgcatttcta aggcgcggct ggaatacctg 1920  
 cgggaccagt tccagattcg tgagaatgac tttcttacct tcgatgccat gcgccacgag 1980  
 gccagtgctg tgggtcgggc catcaggggc aagacggact acggcctcat ggtctttgcc 2040  
 gacaagcggc ttgcccgtgg ggacaagcgg gggaagctgc cccgctggat ccaggagcac 2100  
 ctacagatg ccaacctcaa cctgaccgtg gacgaggggtg tccaggtggc caagtacttc 2160  
 ctgcggcaga tggcacagcc cttccaccgg gaggatcagc tgggcctgtc cctgctcagc 2220  
 ctggagcagc tagaatcaga ggagacgctg aagaggatag agcagattgc tcagcagctc 2280  
 tgagtggggc ggggtggggc ataaacgggt cctggtga 2318

<210> 4  
 <211> 760  
 <212> PRT  
 <213> Homo sapiens

<400> 4

Met Lys Leu Asn Val Asp Gly Leu Leu Val Tyr Phe Pro Tyr Asp Tyr  
 1 5 10 15

Ile Tyr Pro Glu Gln Phe Ser Tyr Met Arg Glu Leu Lys Arg Thr Leu  
 20 25 30

Asp Ala Lys Gly His Gly Val Leu Glu Met Pro Ser Gly Thr Gly Lys  
 35 40 45

Thr Val Ser Leu Leu Ala Leu Ile Met Ala Tyr Gln Arg Ala Tyr Pro  
 50 55 60

Leu Glu Val Thr Lys Leu Ile Tyr Cys Ser Arg Thr Val Pro Glu Ile

|   |                         |                     |             |
|---|-------------------------|---------------------|-------------|
| 65  | 70                      | 75                  | 80          |
| Glu Lys Val Ile   | Glu Glu Leu Arg Lys     | Leu Leu Asn Phe Tyr | Glu Lys     |
|   | 85                      | 90                  | 95          |
| Gln Glu Gly   | Glu Lys Leu Pro Phe     | Leu Gly Leu Ala Leu | Ser Ser Arg |
|   | 100                     | 105                 | 110         |
| Lys Asn Leu Cys Ile His Pro                                     | Glu Val Thr Pro Leu Arg | Phe Gly Lys         |             |
|   | 115                     | 120                 | 125         |
| Asp Val Asp Gly Lys Cys His Ser Leu Thr Ala Ser Tyr Val Arg Ala |                         |                     |             |
|   | 130                     | 135                 | 140         |
| Gln Tyr Gln His Asp Thr Ser Leu Pro His Cys Arg Phe Tyr Glu Glu |                         |                     |             |
|   | 145                     | 150                 | 155         |
| Phe Asp Ala His Gly Arg Glu Val Pro Leu Pro Ala Gly Ile Tyr Asn |                         |                     |             |
|   | 165                     | 170                 | 175         |
| Leu Asp Asp Leu Lys Ala Leu Gly Arg Arg Gln Gly Trp Cys Pro Tyr |                         |                     |             |
|   | 180                     | 185                 | 190         |
| Phe Leu Ala Arg Tyr Ser Ile Leu His Ala Asn Val Val Val Tyr Ser |                         |                     |             |
|   | 195                     | 200                 | 205         |
| Tyr His Tyr Leu Leu Asp Pro Lys Ile Ala Asp Leu Val Ser Lys Glu |                         |                     |             |
|   | 210                     | 215                 | 220         |
| Leu Ala Arg Lys Ala Val Val Val Phe Asp Glu Ala His Asn Ile Asp |                         |                     |             |
|   | 225                     | 230                 | 235         |
| Asn Val Cys Ile Asp Ser Met Ser Val Asn Leu Thr Arg Arg Thr Leu |                         |                     |             |
|   | 245                     | 250                 | 255         |
| Asp Arg Cys Gln Gly Asn Leu Glu Thr Leu Gln Lys Thr Val Leu Arg |                         |                     |             |
|   | 260                     | 265                 | 270         |
| Ile Lys Glu Thr Asp Glu Gln Arg Leu Arg Asp Glu Tyr Arg Arg Leu |                         |                     |             |
|   | 275                     | 280                 | 285         |
| Val Glu Gly Leu Arg Glu Ala Ser Ala Ala Arg Glu Thr Asp Ala His |                         |                     |             |
|   | 290                     | 295                 | 300         |
| Leu Ala Asn Pro Val Leu Pro Asp Glu Val Leu Gln Glu Ala Val Pro |                         |                     |             |
|   | 305                     | 310                 | 315         |
| Gly Ser Ile Arg Thr Ala Glu His Phe Leu Gly Phe Leu Arg Arg Leu |                         |                     |             |
|   | 325                     | 330                 | 335         |
| Leu Glu Tyr Val Lys Trp Arg Leu Arg Val Gln His Val Val Gln Glu |                         |                     |             |
|   | 340                     | 345                 | 350         |

Ser Pro Pro Ala Phe Leu Ser Gly Leu Ala Gln Arg Val Cys Ile Gln  
 355 360 365  
 Arg Lys Pro Leu Arg Phe Cys Ala Glu Arg Leu Arg Ser Leu Leu His  
 370 375 380  
 Thr Leu Glu Ile Thr Asp Leu Ala Asp Phe Ser Pro Leu Thr Leu Leu  
 385 390 395 400  
 Ala Asn Phe Ala Thr Leu Val Ser Thr Tyr Ala Lys Gly Phe Thr Ile  
 405 410 415  
 Ile Ile Glu Pro Phe Asp Asp Arg Thr Pro Thr Ile Ala Asn Pro Ile  
 420 425 430  
 Leu His Phe Ser Cys Met Asp Ala Ser Leu Ala Ile Lys Pro Val Phe  
 435 440 445  
 Glu Arg Phe Gln Ser Val Ile Ile Thr Ser Gly Thr Leu Ser Pro Leu  
 450 455 460  
 Asp Ile Tyr Pro Lys Ile Leu Asp Phe His Pro Val Thr Met Ala Thr  
 465 470 475 480  
 Phe Thr Met Thr Leu Ala Arg Val Cys Leu Cys Pro Met Ile Ile Gly  
 485 490 495  
 Arg Gly Asn Asp Gln Val Ala Ile Ser Ser Lys Phe Glu Thr Arg Glu  
 500 505 510  
 Asp Ile Ala Val Ile Arg Asn Tyr Gly Asn Leu Leu Leu Glu Met Ser  
 515 520 525  
 Ala Val Val Pro Asp Gly Ile Val Ala Phe Phe Thr Ser Tyr Gln Tyr  
 530 535 540  
 Met Glu Ser Thr Val Ala Ser Trp Tyr Glu Gln Gly Ile Leu Glu Asn  
 545 550 555 560  
 Ile Gln Arg Asn Lys Leu Leu Phe Ile Glu Thr Gln Asp Gly Ala Glu  
 565 570 575  
 Thr Ser Val Ala Leu Glu Lys Tyr Gln Glu Ala Cys Glu Asn Gly Arg  
 580 585 590  
 Gly Ala Ile Leu Leu Ser Val Ala Arg Gly Lys Val Ser Glu Gly Ile  
 595 600 605  
 Asp Phe Val His His Tyr Gly Arg Ala Val Ile Met Phe Gly Val Pro  
 610 615 620  
 Tyr Val Tyr Thr Gln Ser Arg Ile Leu Lys Ala Arg Leu Glu Tyr Leu  
 625 630 635 640

Arg Asp Gln Phe Gln Ile Arg Glu Asn Asp Phe Leu Thr Phe Asp Ala  
645 650 655

Met Arg His Ala Ala Gln Cys Val Gly Arg Ala Ile Arg Gly Lys Thr  
660 665 670

Asp Tyr Gly Leu Met Val Phe Ala Asp Lys Arg Phe Ala Arg Gly Asp  
675 680 685

Lys Arg Gly Lys Leu Pro Arg Trp Ile Gln Glu His Leu Thr Asp Ala  
690 695 700

Asn Leu Asn Leu Thr Val Asp Glu Gly Val Gln Val Ala Lys Tyr Phe  
705 710 715 720

Leu Arg Gln Met Ala Gln Pro Phe His Arg Glu Asp Gln Leu Gly Leu  
725 730 735

Ser Leu Leu Ser Leu Glu Gln Leu Glu Ser Glu Glu Thr Leu Lys Arg  
740 745 750

Ile Glu Gln Ile Ala Gln Gln Leu  
755 760

<210> 5

<211> 9731

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthesized retroviral vectors

<400> 5

|  |     |
|--|-----|
| caggtggcac ttttcgggga aatgtgcgcg gaacccttat ttgtttat   | 60  |
| ttctaaatac attcaaata                                   |     |
| gtatccgctc atgagacaat aaccctgata aatgcttcaa taatattgaa | 120 |
| aaaggaagag tatgagtatt caacatttcc gtgtcgccct tattcccttt | 180 |
| tttgccttcc tgtttttgct caccagaaa cgctggtgaa agtaaaagat  | 240 |
| gctgaagatc agttgggtgc acgagtgggt tacatcgaac tggatctcaa | 300 |
| cagcggtaag atccttgaga gttttcgccc cgaagaacgt            | 360 |
| tttccaatga tgagcacttt taaagtctctg ctatgtggcg           | 420 |
| cggtattatc ccgtattgac gccgggcaag agcaactcgg            | 480 |
| tcgccgata cactattctc agaatgactt ggttgagtac             | 540 |
| tcaccagtca cagaaaagca tcttacggat ggcattgacag           | 600 |
| taagagaatt atgcagtgtc gccataacca tgagtataa             | 660 |
| cactgcgcc aacttacttc tgacaacgat cggaggaccg             | 720 |
| aaggagctaa ccgctttttt gcacaacatg ggggatcatg            | 780 |
| taactcgcct tgatcggttg gaaccggagc tgaatgaag             | 840 |
| cataccaaac gacgagcgtg acaccacgat gcctgtagca            | 900 |
| atggcaacaa cgttgcgcaa actattaact ggcgaaactac           |     |
| ttactctagc ttcccggcaa caattaatag actggatgga            |     |
| ggcggataaa gttgcaggac cacttctgcg                       |     |
| ctcgccctt ccggctggct ggtttattgc                        |     |
| tgataaatct ggagccggtg agcgtgggtc                       |     |
| tcgcggtatc attgcagcac tggggccaga                       |     |
| tggttaagccc tcccgtatcg                                 |     |

|            |            |            |             |            |             |      |
|------------|------------|------------|-------------|------------|-------------|------|
| tagttatcta | cacgacggg  | agtcaggcaa | ctatggatga  | acgaaataga | cagatcgctg  | 960  |
| agataggtgc | ctcactgatt | aagcattggt | aactgtcaga  | ccaagtttac | tcatatatac  | 1020 |
| tttagattga | tttaaaactt | catttttaat | ttaaaaggat  | ctaggtgaag | atcctttttg  | 1080 |
| ataatctcat | gaccaaatac | ccttaacgtg | agttttcggt  | ccactgagcg | tcagaccccc  | 1140 |
| tagaaaagat | caaaggatct | tcttgagatc | ctttttttct  | gcgcgtaatc | tgctgcttgc  | 1200 |
| aaacaaaaaa | accaccgcta | ccagcgggtg | tttgtttgcc  | ggatcaagag | ctaccaactc  | 1260 |
| tttttccgaa | ggtaactggc | ttcagcagag | cgcagatacc  | aaatactgtc | cttctagtgt  | 1320 |
| agccgtagtt | aggccaccac | ttcaagaact | ctgtagcacc  | gcctacatac | ctcgtctctg  | 1380 |
| taatcctgtt | accagtggct | gctgccagtg | gcgataagtc  | gtgtcttacc | gggttggtact | 1440 |
| caagacgata | gttaccggat | aaggcgcagc | ggtcgggctg  | aacggggggt | tcgtgcacac  | 1500 |
| agcccagctt | ggagcgaacg | acctacaccg | aactgagata  | cctacagcgt | gagctatgag  | 1560 |
| aaagcgccac | gcttcccgaa | gggagaaagg | cggacaggta  | tccggtaagc | ggcagggtcg  | 1620 |
| gaacaggaga | gcgcacgagg | gagcttccag | ggggaaacgc  | ctggtatctt | tatagtcctg  | 1680 |
| tcgggtttcg | ccacctctga | cttgagcgtc | gattttttgtg | atgctcgtca | ggggggcgga  | 1740 |
| gcctatggaa | aaacgccagc | aacgcggcct | ttttacgggt  | cctggccttt | tgctggcctt  | 1800 |
| ttgctcacat | gttctttcct | gcgttatccc | ctgattctgt  | ggataaccgt | attaccgcct  | 1860 |
| ttgagtgagc | tgataccgct | cgccgcagcc | gaacgaccga  | gcgcagcgag | tcagtgagcg  | 1920 |
| aggaagcgga | agagcgccca | atacgcaaac | cgcctctccc  | cgcgcgttgg | ccgattcatt  | 1980 |
| aatgcagctg | gcacgacagg | tttcccgact | ggaaagcggg  | cagtgagcgc | aacgcaatta  | 2040 |
| atgtgagtta | gctcactcat | taggcacccc | aggctttaca  | ctttatgctt | ccggctcgta  | 2100 |
| tggtgtgtgg | aattgtgagc | ggataacaat | ttcacacagg  | aaacagctat | gaccatgatt  | 2160 |
| acgccaagcg | cgcaattaac | cctcactaaa | gggaacaaaa  | gctggagctg | caagcttaat  | 2220 |
| gtagtcttat | gcaatactct | tgtagtcttg | caacatggta  | acgatgagtt | agcaacatgc  | 2280 |
| cttacaagga | gagaaaaagc | accgtgcatg | ccgattgggtg | gaagtaaggt | ggtacgatcg  | 2340 |
| tgccttatta | ggaaggcaac | agacgggtct | gacatggatt  | ggacgaacca | ctgaattgcc  | 2400 |
| gcattgcaga | gatattgtat | ttaagtgcct | agctcgatac  | aataaacggg | tctctctggt  | 2460 |
| tagaccagat | ctgagcctgg | gagctctctg | gctaactagg  | gaaccactg  | cttaagcctc  | 2520 |
| aataaagctt | gccttgagtg | cttcaagtag | tgtgtgcccg  | tctgttgtgt | gactctggta  | 2580 |
| actagagatc | cctcagaccc | ttttagtcag | tgtggaaaat  | ctctagcagt | ggcgcccgaa  | 2640 |
| cagggacctg | aaagcgaaa  | ggaaaccaga | gctctctcga  | cgcaggactc | ggcttgctga  | 2700 |
| agcgcgcacg | gcaagaggcg | agggggcgcg | actggtgagt  | acgcaaaaa  | ttttgactag  | 2760 |
| cggaggctag | aaggagagag | atgggtgcga | gagcgtcagt  | attaagcggg | ggagaattag  | 2820 |
| atcgcgatgg | gaaaaaatc  | ggttaaggcc | agggggaaag  | aaaaaatata | aattaaaaca  | 2880 |
| tatagtatgg | gcaagcaggg | agctagaacg | attcgagtt   | aatcctggcc | tgtagaaac   | 2940 |
| atcagaaggc | tgtagacaaa | tactgggaca | gctacaacca  | tcccttcaga | caggatcaga  | 3000 |

|  |      |
|--|------|
| agaacttaga tcattatata atacagtagc aaccctctat tgtgtgcatc aaaggataga  | 3060 |
| gataaaagac accaaggaag ctttagacaa gatagaggaa gagcaaaaca aaagtaagac  | 3120 |
| caccgcacag caagcggccg ctgatcttca gacctggagc gctcgaggcg acttacctct  | 3180 |
| ctagagtcgg tgtcttctat ggaggtcaaa acagcgtgga tggcgtctcc aggcgatctg  | 3240 |
| acggttctact aaacgagctc tgcttatata gacctccac cgtacacgcc taccgcccac  | 3300 |
| ttgcgtcaat gggcgaggat tgttacgaca ttttgaaaag tcccgttgat tttggtgcca  | 3360 |
| aaacaaactc ccattgacgt caatggggtg gagacttgga aatccccgtg agtcaaaccg  | 3420 |
| ctatccacgc ccattgatgt actgccaaaa ccgcatcacc atggtaatag cgtatgactaa | 3480 |
| tacgtagatg tactgccaaag taggaaagtc ccataaggtc atgtactggg cataatgcca | 3540 |
| ggcgggcat ttaccgtcat tgacgtcaat agggggcgta cttggcatat gatacacttg   | 3600 |
| atgtactgcc aagtgggcag tttaccgtaa atactccacc cattgacgtc aatggaaagt  | 3660 |
| ccctattggc gttactatgg gaacatacgt cattattgac gtcaatgggc gggggtcgtt  | 3720 |
| ggcggtcag ccaggcgggc catttaccgt aagttatgta acgcggaact cccaagctta   | 3780 |
| tcgaggagga gatatgaggg acaattggag aagtgaatta tataaatata aagtagtaaa  | 3840 |
| aattgaacca ttaggagtag caccaccaa ggcaaagaga agagtgggtc agagagaaaa   | 3900 |
| aagagcagtg ggaataggag ctttgttcct tgggttcttg ggagcagcag gaagcactat  | 3960 |
| ggcgcgagcc tcaatgacgc tgacggtaca ggccagacaa ttattgtctg gtatagtga   | 4020 |
| gcagcagaac aatttgctga gggctattga ggcgcaacag catctgttg aactcacagt   | 4080 |
| ctggggcatc aagcagctcc aggcaagaat cctggctgtg gaaagatacc taaaggatca  | 4140 |
| acagctcctg gggatttggg gttgctctgg aaaactcatt tgcaccactg ctgtgccttg  | 4200 |
| gaatgctagt tggagtaata aatctctgga acagattgga atcacacgac ctggatggag  | 4260 |
| tgggacagag aaattaacaa ttacacaagc ttaatacact ccttaattga agaatcgcaa  | 4320 |
| aaccagcaag aaaagaatga acaagaatta ttggaattag ataaatgggc aagtttgtgg  | 4380 |
| aattggttta acataacaaa ttggctgtgg tatataaaat tattcataat gatagtagga  | 4440 |
| ggcttggtag gtttaagaat agtttttctg gtactttcta tagtgaatag agttaggcag  | 4500 |
| ggatattcac cattatcggt tcagaccac ctccaaccc cgaggggacc cgacaggccc    | 4560 |
| gaaggaatag aagaagaagg tggagagaga gacagagaca gatccattcg attagtgaac  | 4620 |
| ggatctcgac ggttaacttt taaaagaaaa ggggggattg gggggtacag tgcaggggaa  | 4680 |
| agaatagtag acataatagc aacagacata caaactaaag aattacaaaa acaaattaca  | 4740 |
| aaaattcaaa attttatcgc atgttctttc ctgcgttacc cctgattct gtggataacc   | 4800 |
| gtattaccgc catgcattag ttattaatag taatcaatta cggggtcatt agttcatagc  | 4860 |
| ccatatatgg agttccgcgt tacataactt acggtaaatg gccgcctgg ctgaccgccc   | 4920 |
| aacgaccccc gccattgac gtcaataatg acgtatgttc ccataagtaac gccaataggg  | 4980 |
| actttccatt gacgtcaatg ggtggagtat ttacggtaaa ctgccactt ggcagtacat   | 5040 |
| caagtgtatc atatgccaag tacgccccct attgacgtca atgacggtaa atggcccgcc  | 5100 |
| tggcattatg ccagtagcat gacctatgg gactttccta cttggcagta catctacgta   | 5160 |

|   |      |
|---|------|
| ttagtcatcg ctattaccat ggtgatgcgg ttttggcagt acatcaatgg gcgtggatag   | 5220 |
| cggtttgact cacggggatt tccaagtctc caccctattg acgtcaatgg gagtttgttt   | 5280 |
| tggcaccaaa atcaacggga ctttccaaaa tgtcgttaaca actccgcccc attgacgcaa  | 5340 |
| atgggcggtg ggcgtgtacg gtgggaggtc tatataagca gagctggttt agtgaaccgt   | 5400 |
| cagatccgct agcgttaccg gactcagatc tgcagctcaa gcttcgaatt ctgcagtcga   | 5460 |
| cgggtaccgag gggccgggat ccaccggtcg ccaccatggc ctctccgag aacgtcatca   | 5520 |
| ccgagttcat gcgcttcaag gtgcgcatgg agggcaccgt gaacggccac gagttcgaga   | 5580 |
| tgcagggcga gggcgagggc cgcccctacg agggccacaa caccgtgaag ctgaagggtga  | 5640 |
| ccaagggcgg cccctgccc ttgcctggg acatcctgtc ccccagttc cagtacggct      | 5700 |
| ccaagggtga cgtgaagcac ccgcccga tccccgacta caagaagctg tccttccccg     | 5760 |
| agggcttcaa gtgggagcgc gtgatgaact tgcaggacgg cggcgtggcg accgtgaccc   | 5820 |
| aggactcctc cctgcaggac ggctgcttca tctacaaggt gaagttcatc ggcgtgaact   | 5880 |
| tcccctccga cggccccgtg atgcagaaga agaccatggg ctgggaggcc tccaccgagc   | 5940 |
| gcctgtaccc ccgcgacggc gtgctgaagg gcgagacca caaggccctg aagctgaagg    | 6000 |
| acggcgggcca ctacctggtg gagttcaagt ccatctacat ggccaagaag ccggtgcagc  | 6060 |
| tgcccggtga ctactacgtg gacgccaagc tggacatcac ctcccacaac gaggactaca   | 6120 |
| ccatcgtgga gcagtacgag cgcaccgagg gccgccacca cctgttcctg tagcggggcc   | 6180 |
| tgcacaatca acctctggat tacaaaattt gtgaaagatt gactggtatt cttaactatg   | 6240 |
| ttgctccttt tacgctatgt ggatacgctg ctttaatgcc tttgtatcat gctattgctt   | 6300 |
| cccgtatggc tttcattttc tcctccttgt ataaatcctg gttgctgtct ctttatgagg   | 6360 |
| agttgtggcc cgttgctcagg caacgtggcg tgggtgtgcac tgtgtttgct gacgcaaccc | 6420 |
| ccactgggtg gggcattgcc accacctgtc agctcctttc cgggactttc gctttcccc    | 6480 |
| tccctattgc cacggcggaa ctcatcgccg cctgccttgc ccgctgctgg acaggggctc   | 6540 |
| ggctgttggg cactgacaat tccgtggtgt tgcggggaa gctgacgtcc tttccatggc    | 6600 |
| tgtcgcctg tgttgccacc tggattctgc gcgggacgtc cttctgctac gtcccttcgg    | 6660 |
| ccctcaatcc agcggacctt ccttcccgcg gcctgctgcc ggctctgcgg cctcttccgc   | 6720 |
| gtcttcgcct tcgccctcag acgagtcgga tctccctttg ggccgcctcc ccgcctggaa   | 6780 |
| ttccgcgact ctagatcata atcagccata ccacatttgt agaggtttta cttgctttaa   | 6840 |
| aaaacctccc acacctcccc ctgaacctga aacataaaat gaatgcaatt gttgttggtta  | 6900 |
| acttgtttat tgcagcttat aatggttaca aataaagcaa tagcatcaca aatttcacaa   | 6960 |
| ataaagcatt tttttcactg cattctagtt gtggtttgtc caaactcatc aatgtatctt   | 7020 |
| aaccaggcgg ggaggcggcc caaagggaga tccgactcgt ctgaggcgga aggcgaagac   | 7080 |
| gcggaagagg ccgcagagcc ggcagcaggc cgcgggaagg aaggtccgct ggattgaggg   | 7140 |
| ccgaaggggac gtagcagaag gacgtcccg cgcagaatcca ggtggcaaca caggcgagca  | 7200 |
| gcatggaaga ggacgtcagc ttccccgaca acaccacgga attgtcagtg cccaacagcc   | 7260 |

|   |      |
|---|------|
| gagccccctgt ccagcagcgg gcaaggcagg cggcgatgag ttccgccgtg gcaataggga  | 7320 |
| gggggaaagc gaaagtcccc gaaaggagct gacaggtggt ggcaatgccc caaccagtgg   | 7380 |
| gggttgcgtc agcaaacaca gtgcacacca cgccacgttg cctgacaacg ggccacaact   | 7440 |
| cctcataaag agacagcaac caggatttat acaaggagga gaaaatgaaa gccatacggg   | 7500 |
| aagcaatagc atgatacaaa ggcattaaag cagcgtatcc acatagcgta aaaggagcaa   | 7560 |
| catagttaag aataccagtc aatctttcac aaattttgta atccagaggt tgattgtcga   | 7620 |
| cgcggccgct ttacttgtac agctcgtcca tgccgagagt gatcccgcg gcggtcacga    | 7680 |
| actocagcag gaccatgtga tcgcgcttct cgttggggtc tttgctcagg gcggactggg   | 7740 |
| tgctcaggta gtggttgtcg ggcagcagca cggggccgtc gccgatgggg gtgttctgct   | 7800 |
| ggtagtggtc ggcgagctgc acgctgccgt cctcgatgtt gtggcggatc ttgaagttca   | 7860 |
| ccttgatgcc gttcttctgc ttgtcggcca tgatatagac gttgtggctg ttgtagttgt   | 7920 |
| actccagctt gtgccccagg atgttgccgt cctccttgaa gtcgatgccc ttcagctcga   | 7980 |
| tgcggttcac cagggtgtcg ccctcgaact tcacctcggc gcgggtcttg tagttgccgt   | 8040 |
| cgtccttgaa gaagatgggt cgctcctgga cgtagccttc gggcatggcg gacttgaaga   | 8100 |
| agtcgtgctg cttcatgtgg tcggggtagc ggctgaagca ctgcacgccg taggtcaggg   | 8160 |
| tggtcacgag ggtggggcag ggcacgggca gcttgccggt ggtgcagatg aacttcaggg   | 8220 |
| tcagcttgcc gtaggtggca tcgccctcgc cctcgccgga cacgctgaac ttgtggccgt   | 8280 |
| ttacgtcgcc gtccagctcg accaggatgg gcaccacccc ggtgaacagc tcctcgccct   | 8340 |
| tgctcaccat ggtggcgacc ggtggatcct gaagaaaagg gagaattcga attcgagctc   | 8400 |
| ggtaccttta agaccaatga cttacaaggc agctgtagat cttagccact ttttaaaga    | 8460 |
| aaagggggga ctggaagggc taattcactc ccaacgaaga caagatctgc tttttgcttg   | 8520 |
| tactgggtct ctctggttag accagatctg agcctgggag ctctctggct aactagggaa   | 8580 |
| cccactgctt aagcctcaat aaagcttgcc ttgagtgcct caagtagtgt gtgcccgctc   | 8640 |
| gttgtgtgac tctggttaact agagatccct cagacccttt tagtcagtgt ggaaaatctc  | 8700 |
| tagcagtagt agttcatgtc atcttattat tcagtattta taacttgcaa agaaatgaat   | 8760 |
| atcagagagt gagaggaaact tgttttattgc agcttataat ggttacaaat aaagcaatag | 8820 |
| catcacaat ttcacaaata aagcattttt ttactgcat tctagttgtg gtttgtcaa      | 8880 |
| actcatcaat gtatcttata atgtctggct ctagctatcc cgcccctaac tccgccagc    | 8940 |
| tccgccatt ctccgcccc tggctgacta atttttttta tttatgcaga ggccgaggcc     | 9000 |
| gcctcggcct ctgagctatt ccagaagtag tgaggaggct tttttggagg cctaggcttt   | 9060 |
| tgcgtcgaga cgtacccaat tcgccctata gtgagtcgta ttacgcgcgc tcaactggccg  | 9120 |
| tcgttttaca acgtcgtgac tgggaaaacc ctggcggttac ccaacttaat cgccttgca   | 9180 |
| cacatcccc tttcgccagc tggcgtaata gcgaagaggc ccgcaccgat cgcccttccc    | 9240 |
| aacagttgcg cagcctgaat ggcgaatggc gcgacgcgcc ctgtagcggc gcattaagcg   | 9300 |
| cggcggtgt ggtggttacg cgcagcgtga ccgctacact tgccagcgcc ctagcgccc     | 9360 |
| ctcctttcgc tttcttccct tcctttctcg ccacgttcgc cggttttccc cgtcaagctc   | 9420 |



|   |      |
|---|------|
| taaatcgggg gctcccttta gggttccgat ttagtgcttt acggcacctc gacccccaaa | 9480 |
| aacttgatta gggatgatgt tcacgtagt ggccatcgcc ctgatagacg gtttttcgcc  | 9540 |
| ctttgacgtt ggagtcacg tcttttaata gtggactctt gttccaaact ggaacaacac  | 9600 |
| tcaaccctat ctcggtctat tcttttgatt tataagggat tttgccgatt tcggcctatt | 9660 |
| ggttaaaaa tgagctgatt taacaaaaat ttaacgcgaa ttttaacaaa atattaacgt  | 9720 |
| ttacaatttc c  | 9731 |

&lt;210&gt; 6

&lt;211&gt; 9782

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Synthesized retroviral vectors

&lt;400&gt; 6

|   |      |
|---|------|
| ccggtcgcca ccatggcctc ctccgagaac gtcacaccg agttcatgcg cttcaagggtg   | 60   |
| cgcattgagg gcaccgtgaa cggccacgag ttccgagatcg agggcgaggg cgagggccgc  | 120  |
| ccctacgagg gccacaacac cgtgaagctg aagggtacca agggcgggcc cctgcccttc   | 180  |
| gcctgggaca tcctgtcccc ccagttccag tacggctcca aggtgtacgt gaagcaccac   | 240  |
| gccgacatcc ccgactacaa gaagctgtcc ttccccgagg gcttcaagtg ggagcgcggtg  | 300  |
| atgaacttcg aggacggcgg cgtggcgacc gtgaccacg actcctccct gcaggacggc    | 360  |
| tgcttcatct acaagggtgaa gttcatcggc gtgaacttcc cctccgacgg ccccgatgatg | 420  |
| cagaagaaga ccatgggctg ggaggcctcc accgagcgcc tgtacccccg cgacggcggtg  | 480  |
| ctgaaggcg agaccacaa ggcctgaag ctgaaggacg gcggccacta cctggtggag      | 540  |
| ttcaagtcca tctacatggc caagaagccc gtgcagctgc ccggctacta ctacgtggac   | 600  |
| gccaagctgg acatcacctc ccacaacgag gactacacca tcgtggagca gtacgagcgc   | 660  |
| accgagggcc gccaccacct gttcctgtag cggccgcgac tctagatcat aatcagccat   | 720  |
| accacatttg tagaggtttt acttgcttta aaaaacctcc cacacctccc cctgaacctg   | 780  |
| aaacataaaa tgaatgcaat tgttgtgttt aacttgttta ttgcagctta taatggttac   | 840  |
| aaataaagca atagcatcac aaatttcaca aataaagcat ttttttact gcattctagt    | 900  |
| tggatccttt atttgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg   | 960  |
| caataaaciaa gtttaacaaca acaattgcat tcattttatg tttcagggtc agggggaggt | 1020 |
| gtgggaggtt ttttaaaagca agtaaaacct ctacaaatgt ggtatggctg aattccaggc  | 1080 |
| ggggaggcgg cccaaaggga gatccgactc gtctgagggc gaaggcgaag acgcggaaga   | 1140 |
| ggccgcagag ccggcagcag gccgcgggaa ggaaggtcog ctggattgag ggccgaaggg   | 1200 |
| acgtagcaga aggacgtccc gcgcagaatc cagggtggca cacaggcgag cagccatgga   | 1260 |
| aaggacgtca gtttccccga caacaccacg gaattgtcag tgcccaacag ccgagcccct   | 1320 |
| gtccagcagc gggcaaggca ggccggcatg agttccgcog tggcaatagg gagggggaaa   | 1380 |
| gcgaaagtcc cggaaaggag ctgacaggtg gtggcaatgc cccaaccagt gggggttgcg   | 1440 |

|  |      |
|--|------|
| tcagcaaaca cagtgcacac cagccacgt tgcctgacaa cgggccacaa ctctcataa    | 1500 |
| agagacagca accaggattt atacaaggag gagaaaatga aagccatacg ggaagcaata  | 1560 |
| gcatgataca aaggcattaa agcagcgtat ccacatagcg taaaaggagc aacatagtta  | 1620 |
| agaataccag tcaatctttc acaaattttg taatccagag gttgattgtc gacgcggccg  | 1680 |
| ctttacttgt acagctcgtc catgccgaga gtgatcccgg cggcggtcac gaactccagc  | 1740 |
| aggaccatgt gatcgcgctt ctctgtgggg tctttgctca gggcggtactg ggtgctcagg | 1800 |
| tagtggttgt cgggcagcag cacggggccg tcgccgatgg ggggtgtctg ctggtagtgg  | 1860 |
| tcggcgagct gcacgctgcc gtcctcgatg ttgtggcgga tcttgaagtt caccttgatg  | 1920 |
| ccgttcttct gcttgctggc catgatatac acgttggtggc tgtttagtgg gtactccagc | 1980 |
| ttgtgcccc ggatgttgcc gtcctccttg aagtcgatgc ccttcagctc gatgcggttc   | 2040 |
| accagggtgt cgccctcgaa cttcacctcg gcgcgggtct tgtagttgcc gtcgtccttg  | 2100 |
| aagaagatgg tgcgctctg gacgtagcct tcgggcatgg cggacttgaa gaagtcgtgc   | 2160 |
| tgcttcatgt ggtcggggta gcggctgaag cactgcacgc cgtaggtcag ggtggtcacg  | 2220 |
| agggtgggcc agggcacggg cagcttgccg gtggtgcaga tgaacttcag ggtcagcttg  | 2280 |
| ccgtaggtgg catcgccctc gccctcgccg gacacgctga acttggtggc gtttacgtcg  | 2340 |
| ccgtccagct cgaccaggat gggcaccacc ccggtgaaca gtcctcgcc cttgctcacc   | 2400 |
| atctgaagaa aaggaggta cctttaagac caatgactta caaggcagct gtagatctta   | 2460 |
| gccacttttt aaaagaaaag gggggactgg aagggtctaat tcaactccaa cgaagacaag | 2520 |
| atatccttga tctgtggatc taccacacac aaggctactt ccctgattgg cagaactaca  | 2580 |
| caccagggcc agggatcaga tatccactga cctttggatg gtgctacaag ctagtaccag  | 2640 |
| ttgagcaaga gaaggtagaa gaagccaatg aaggagagaa caccgccttg ttacaccctg  | 2700 |
| tgagcctgca tgggatggat gaccgggaga gagaagtatt agagtggagg tttgacagcc  | 2760 |
| gcctagcatt tcatcacatg gcccgagagc tgcacccgga gtacttcaag aactgctgac  | 2820 |
| atcgagcttg ctacaaggga ctttccgctg gggactttcc agggaggcgt ggctgggcg   | 2880 |
| ggactgggga gtggcgagcc ctacagatgct gcatataagc agctgctttt tgcttgact  | 2940 |
| gggtctctct ggttagacca gatctgagcc tgggagctct ctggctaact agggaacca   | 3000 |
| ctgcttaagc ctcaataaag cttgccttga gtgcttcaag tagtgtgtgc ccgtctgttg  | 3060 |
| tgtgactctg gtaactagag atccctcaga cccttttagt cagtgtggaa aatctctagc  | 3120 |
| agtagtagtt catgtcatct tattattcag tatttataac ttgcaaagaa atgaatatca  | 3180 |
| gagagtgaga ggccttgaca ttataataga tttagcagga attgaactag gagtggagca  | 3240 |
| cacaggcaaa gctgcagaag tacttggaag aagccaccag agatactcac gattctgcac  | 3300 |
| atacctggct aatcccagat cctaaggatt acattaagtt tactaacatt tatataatga  | 3360 |
| tttatagttt aaagtataaa cttatctaatt ttactattct gacagatatt aattaatcct | 3420 |
| caaatatcat aagagatgat tactattatc ccattttaac acaagaggaa actgagaggg  | 3480 |
| aaagatgttg aagtaatttt cccacaatta cagcatccgt tagttacgac tctatgatct  | 3540 |
| tctgacacaa attccattta ctctcacc cttatgactcag tcgaatatat caaagttatg  | 3600 |

|  |      |
|--|------|
| gacattatgc taagtaacaa attacccttt tatatagtaa atactgagta gattgagaga  | 3660 |
| agaaattgtt tggcaaacct gaatagcttc cagaagaaga gaagtgagga taagaataac  | 3720 |
| agttgtcatt aaccagtttt aacaagtaac ttggttagaa agggattcaa atgcataaag  | 3780 |
| caagggataa atttttcttg caacaagact atacaatata accttaaata tgacttcaaa  | 3840 |
| taattgttgg aacttgataa aactaattaa atattattga agattatcaa tattataaat  | 3900 |
| gtaatttact tttaaaaagg gaacatagaa atgtgtatca ttagagtaga aaacaatcct  | 3960 |
| tattatcaca atttgtcaaa acaagtttgt tattaacaca agtagaatac tgcattcaat  | 4020 |
| taagttgact gcagattttg tgttttggtt aaattagaaa gagataacaa caatttgaat  | 4080 |
| tattgaaagt aacatgtaaa tagttctaca tacgttcttt tgacatcttg ttcaatcatt  | 4140 |
| gatcgaagtt ctttatcttg gaagaatttg ttccaaagac tctgaaataa ggaaaacaat  | 4200 |
| ctattatata gtctcacacc tttgttttac ttttagtgat ttcaatttaa taatgtaaat  | 4260 |
| ggttaaaatt tattcttctc tgagatcatt tcacattgca gatagaaaac ctgagactgg  | 4320 |
| ggtaattttt attaaaatct aatttaatct cagaacacaca tctttattct aacatcaatt | 4380 |
| tttccagttt gatattatca tataaagtca gccttcctca tctgcaggtt ccacaacaaa  | 4440 |
| aatccaacca actgtggatc aaaaatattg ggaaaaaatt aaaaatagca atacaacaat  | 4500 |
| aaaaaaatac aaatcagaaa aacagcacag tataacaact ttatttagca ttacaatct   | 4560 |
| attaggtatt ataagtaatc tagaattaat tccgtgtatt ctatagtgtc acctaaatcg  | 4620 |
| tatgtgtatg atacataagg ttatgtatta attgtagccg cgttctaacg acaatatgta  | 4680 |
| caagcctaatt tgtgtagcat ctggcttact gaagcagacc ctatcatctc tctcgtaaac | 4740 |
| tgccgtcaga gtcggtttgg ttggacgaac cttctgagtt tctggtaacg ccgttcgca   | 4800 |
| ccccggaat ggtcagcgaa ccaatcagca gggcatcgc tagccagatc ctctacgccg    | 4860 |
| gacgcacgtt ggccggcatc accggcgcca caggtgcggt tgctggcgcc tatatcgccg  | 4920 |
| acatcaccga tggggaagat cgggctcgcc acttcgggct catgagcgct tgtttcggcg  | 4980 |
| tgggtatggt ggcaggcccc gtggccgggg gactgttggg cgccatctcc ttgcatgcac  | 5040 |
| cattccttgc ggcggcggtg ctcaacggcc tcaacctact actgggctgc ttctaatgc   | 5100 |
| aggagtgcga taaggagag cgtcgatatg gtgcactctc agtacaatct gctctgatgc   | 5160 |
| cgcatagtta agccagcccc gacaccgcc aacaccgct gacgcgcct gacgggcttg     | 5220 |
| tctgtcccg gcatccgctt acagacaagc tgtgaccgtc tccgggagct gcatgtgtca   | 5280 |
| gaggttttca ccgtcatcac cgaaacgcgc gagacgaaag ggcctcgtga tacgcctatt  | 5340 |
| tttatagggt aatgtcatga taataatggt ttcttagacg tcaggtggca cttttcgggg  | 5400 |
| aaatgtgcgc ggaacccta tttgtttatt tttctaaata cattcaaata tgtatccgct   | 5460 |
| catgagacaa taaccctgat aaatgcttca ataattattga aaaaggaaga gtatgagtat | 5520 |
| tcaacatttc cgtgtcgccc ttattccctt ttttgcggca ttttgccttc ctgtttttgc  | 5580 |
| tcaccagaa acgctgggtg aagtaaaaga tgctgaagat cagttgggtg cacgagtggg   | 5640 |
| ttacatcgaa ctggatctca acagcggtaa gatccttgag agttttcgcc ccgaagaacg  | 5700 |

|   |      |
|---|------|
| ttttccaatg atgagcactt ttaaagttct gctatgtggc gcggtattat cccgtattga | 5760 |
| cgccgggcaa gagcaactcg gtcgccgcat acactattct cagaatgact tggttgagta | 5820 |
| ctcaccagtc acagaaaagc atcttacgga tggcatgaca gtaagagaat tatgcagtgc | 5880 |
| tgccataacc atgagtgata aactgcggc caacttactt ctgacaacga tcggaggacc  | 5940 |
| gaaggagcta accgcttttt tgcacaacat gggggatcat gtaactcgcc ttgatcgttg | 6000 |
| ggaaccggag ctgaatgaag ccataccaaa cgacgagcgt gacaccacga tgcctgtagc | 6060 |
| aatggcaaca acgttgcgca aactattaac tggcgaacta cttactctag cttcccgga  | 6120 |
| acaattaata gactggatgg aggcggataa agttgcagga ccacttctgc gctcggccct | 6180 |
| tccggctggc tggttttattg ctgataaatc tggagccggt gagcgtgggt ctgcggtat | 6240 |
| cattgcagca ctggggccag atggtaagcc ctcccgtatc gtagttatct acacgacggg | 6300 |
| gagtcaggca actatggatg aacgaaatag acagatcgct gagatagggt cctcactgat | 6360 |
| taagcattgg taactgtcag accaagttta ctcatatata ctttagattg atttaaaact | 6420 |
| tcatttttaa tttaaaagga tctaggtgaa gatccttttt gataatctca tgacaaaat  | 6480 |
| cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga tcaaaggatc | 6540 |
| ttcttgagat cttttttttc tgcgcgtaat ctgctgcttg caaacaacaa aaccaccgt  | 6600 |
| accagcggtg gtttgtttgc cggatcaaga gctaccaact ctttttccga aggtaactgg | 6660 |
| cttcagcaga gcgcagatac caaatactgt ccttctagt tagccgtagt taggccacca  | 6720 |
| cttcaagaac tctgtagcac cgcctacata cctcgctctg ctaatcctgt taccagtggc | 6780 |
| tgctgccagt ggcgataagt cgtgtcttac cgggttgac tcaagacgat agttaccgga  | 6840 |
| taaggcgag cggtcgggct gaacgggggg ttcgtgcaca cagcccagct tggagcgaac  | 6900 |
| gacctacacc gaactgagat acctacagcg tgagctatga gaaagcgcca cgcttccga  | 6960 |
| aggagaaaag gcggacaggt atccggtgag cggcagggtc ggaacaggag agcgcacgag | 7020 |
| ggagcttcca gggggaaacg cctggtatct ttatagtcct gtcgggtttc gccacctctg | 7080 |
| acttgagcgt cgatttttgt gatgctcgtc aggggggagg agcctatgga aaaacgccag | 7140 |
| caacgcggcc tttttacggt tcctggcctt ttgctggcct ttgctcaca tgttctttcc  | 7200 |
| tgcttatcc cctgattctg tggataaccg tattaccgcc tttgagttag ctgataccgc  | 7260 |
| tcgccgcagc cgaacgaccg agcgacgca gtcagtgagc gaggaagcgg aagagcgccc  | 7320 |
| aatacgcaaa ccgcctctcc ccgcgcgttg gccgattcat taatgcagct gtggaatgtg | 7380 |
| tgtcagttag ggtgtggaat gtccccaggc tccccagcag gcagaagtat gcaaagcatg | 7440 |
| catctcaatt agtcagcaac caggtgtgga aagtccccag gctccccagc aggcagaagt | 7500 |
| atgcaaagca tgcatctcaa ttagtcagca accatagtcc cgcccctaac tccgccatc  | 7560 |
| ccgccctaa ctccgccag ttccgccat tctccgccc atggctgact aatttttttt     | 7620 |
| atztatgcag aggcggaggc cgctcgcc tctgagctat tccagaagta gtgaggaggc   | 7680 |
| ttttttggag gcctaggctt ttgcaaaaag cttggacaca agacaggctt gcgagatatg | 7740 |
| tttgagaata ccactttatc ccgcgtcagg gagaggcagt gcgtaaaaag acgcggactc | 7800 |
| atgtgaaata ctggttttta gtgcgccaga tctctataat ctgcgcgaac ctattttccc | 7860 |

|             |             |             |            |            |             |      |
|-------------|-------------|-------------|------------|------------|-------------|------|
| ctcgaacact  | ttttaagccg  | tagataaaca  | ggctgggaca | cttcacatga | gcgaaaaata  | 7920 |
| catcgtcacc  | tgggacatgt  | tgcagatcca  | tgcacgtaaa | ctcgcaagcc | gactgatgcc  | 7980 |
| ttctgaacaa  | tggaaggca   | ttattgccgt  | aagccgtggc | ggtctgtacc | gggtgcgtta  | 8040 |
| ctggcgcgtg  | aactgggtat  | tcgtcatgtc  | gataccgttt | gtatttccag | ctacgatcac  | 8100 |
| gacaaccagc  | gcgagcttaa  | agtgtgaaa   | cgcgcagaag | gcgatggcga | aggcttcac   | 8160 |
| gttattgatg  | acctgggtga  | taccgggtgt  | actgcggttg | cgattcgtga | aatgtatcca  | 8220 |
| aaagcgcaact | ttgtcaccat  | cttcgcaaaa  | ccggctggtc | gtccgctggt | tgatgactat  | 8280 |
| gttggtgata  | tcccgcaga   | tacctggatt  | gaacagccgt | gggatatggg | cgtcgtattc  | 8340 |
| gtcccgcgcaa | tctccggtcg  | ctaattcttt  | caacgcctgg | cactgccggg | cgttgttctt  | 8400 |
| tttaacttca  | ggcgggttac  | aatagtttcc  | agtaagtatt | ctggaggctg | catccatgac  | 8460 |
| acaggcaaac  | ctgagcgaaa  | ccctgttcaa  | accccgcttt | aaacatcctg | aaacctcgac  | 8520 |
| gctagtcogc  | cgctttaatc  | acggcgacac  | accgcctgtg | cagtcggccc | ttgatggtaa  | 8580 |
| aaccatccct  | cactgggtatc | gcatgattaa  | ccgtctgatg | tggatctggc | gcggcattga  | 8640 |
| cccacgcgaa  | atcctcgacg  | tccaggcacg  | tattgtgatg | agcgatgccg | aacgtaccga  | 8700 |
| cgatgattta  | tacgatacgg  | tgattggcta  | ccgtggcggc | aactggattt | atgagtgggc  | 8760 |
| cccggtatctt | tgtgaaggaa  | ccttacttct  | gtggtgtgac | ataattggac | aaactaccta  | 8820 |
| cagagattta  | aagctctaag  | gtaaatataa  | aatttttaag | tgtataatgt | gttaaactac  | 8880 |
| tgatttcta   | tgtttgtgta  | ttttagattc  | caacctatgg | aactgatgaa | tgaggagcagt | 8940 |
| ggtggaatgc  | ctttaatgag  | gaaaacctgt  | tttgctcaga | agaaatgcc  | tctagtgatg  | 9000 |
| atgaggctac  | tgctgactct  | caacattcta  | ctcctccaaa | aaagaagaga | aaggtagaag  | 9060 |
| acccaagga   | ctttccttca  | gaattgctaa  | gttttttgag | tcatgctgtg | tttagtaata  | 9120 |
| gaactcttgc  | ttgctttgct  | atttacacca  | caaaggaaaa | agctgcactg | ctatacaaga  | 9180 |
| aaattatgga  | aaaatattct  | gtaaccttta  | taagtaggca | taacagttat | aatcataaca  | 9240 |
| tactgttttt  | tcttactcca  | cacaggcata  | gagtgtctgc | tattaataac | tatgctcaaa  | 9300 |
| aattgtgtac  | ctttagcttt  | ttaatttgta  | aaggggttaa | taaggaatat | ttgatgtata  | 9360 |
| gtgccttgac  | tagagatcat  | aatcagccat  | accacatttg | tagaggtttt | acttgcttta  | 9420 |
| aaaaacctcc  | cacacctccc  | cctgaacctg  | aaacataaaa | tgaatgcaat | tggtgtgtgt  | 9480 |
| aacttgttta  | ttgcagctta  | taatgggttac | aaataaagca | atagcatcac | aaatttcaca  | 9540 |
| aataaagcat  | ttttttcact  | gcattctagt  | tgtgggttgt | ccaaactcat | caatgtatct  | 9600 |
| tatcatgtct  | ggatcaactg  | gataactcaa  | gctaacccaa | atcatcccaa | acttcccacc  | 9660 |
| ccatacccta  | ttaccactgc  | caattacctg  | tggtttcatt | tactctaaac | ctgtgattcc  | 9720 |
| tctgaattat  | tttcatttta  | aagaaattgt  | atttggttaa | tatgtactac | aaacttagta  | 9780 |
| gt          |             |             |            |            |             | 9782 |

**THIS PAGE BLANK (USPTO)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**